

**Southern
Region**



**Drought
Ready
Tasmania**

**Drought Risk,
Resilience &
Adaptive
Capacity Data**



RMCG


Australian Government
Department of Agriculture,
Fisheries and Forestry



**Future
Drought
Fund**


**Tasmanian
Government**

Contents

Executive Summary	1
Purpose	1
The Southern Region	1
Key Findings.....	1
Recommendations	4
Glossary.....	5
1. Introduction.....	6
1.1. Assessing the impacts of drought	6
1.2. Project Purpose	6
1.3. Definition of drought	6
1.4. Assessment framework	7
1.5. Quantitative analysis	10
2. Regional Profile.....	11
2.1. Overview	11
2.2. Agriculture and Industry.....	12
2.3. Natural Environment	16
2.4. Community	18
3. Past and Current Impacts of Drought.....	21
3.1. Past Climate Trends for Key Townships .	23
3.2. Impacts on Agriculture	28
3.3. Impacts on the Natural Environment.....	29
3.4. Impacts on Communities.....	30
4. Future Impacts of Drought.....	32
4.1. Future Drought Conditions	32
4.2. Future Climate Trends for the Southern Region	32
Water Availability	35

4.3.	Impacts on Agriculture	36
4.4.	Impacts on Natural Environment	40
4.5.	Impacts on Communities	42
5.	Risk, Adaptive Capacity & Resilience	43
5.1.	Drought Impact (Risk)	43
5.2.	Adaptive Capacity	44
5.3.	Resilience to Drought.....	49
5.4.	Gaps in Preparedness	50
6.	Building Resilience to Drought	52
6.1.	Key Observations and Themes	52
6.2.	Recommendations	53
	References	54
	Appendix 1: Adaptive capacity and vulnerability assessment methodology	57
1	Introduction.....	57
2	Determining Adaptive Capacity.....	57
2.1	Physical Capital	57
2.2	Natural Capital	58
2.3	Financial Capital	58
2.4	Human Capital	58
2.5	Social Capital	58
2.6	Combining the Measures	58
3	Determining Drought Impact (Risk)	59
4	Determining Drought Vulnerability & Resilience	59
	Appendix 2: Socio-economic data (SEIFA)	60
	Appendix 3: Regional economic data – income, housing, education	63

Executive Summary

Purpose

The purpose of this report is to provide a snapshot in time of the indicative and potential drought impacts for the Southern region of Tasmania. It answers the following three questions:

1. What is the prevalence, severity and impacts of drought? (Past/current)
2. What is the likely prevalence, severity and impact of droughts? (Future)
3. What are the vulnerabilities, gaps in preparedness and adaptive capacity for drought and other related permanent transitions to a changing climate? (Analysis)

The Southern region drought data report will support the engagement activities of the Regional Project Coordinator and underpin the development of the Regional Drought Resilience Plan for the Southern region with regional stakeholders.

Drought resilience is the ability to adapt, reorganise or transform in response to changing temperature, increasing variability and scarcity of rainfall and changed seasonality of rainfall, for improved economic, environmental and social wellbeing. This report analyses the resilience of agricultural, natural environment and community systems to drought.

The Southern Region

The Southern region has a land area of 25,964 km², a population of just under 300,000 and is comprised of the following Local Government Areas (LGAs): Brighton, Central Highlands, Clarence, Derwent Valley, Glamorgan-Spring Bay, Glenorchy, Hobart, Huon Valley, Kingborough, Sorell, Southern Midlands and Tasman.

Compared to Tasmania as a whole, households in the Southern region are skewed towards the higher end of household incomes, people tend to be more educated and slightly younger. Clarence, Hobart and Kingborough have the least level of disadvantage of the LGAs in the Southern region, while people within the Brighton and Central Highlands LGA demonstrate the most disadvantage.

Only 3.5% of the population in the Southern region are employed in agriculture, forestry and fisheries. However, this increases in some LGAs, such as Huon Valley and the Southern Midlands where these sectors employ around a quarter of the population.

Huon Valley and Southern Midlands also account for around 41% of the region's agricultural output. Fruit (including nuts) is the region's largest contributor to the value of agriculture, this is followed by livestock products (excluding dairy), and field crops (including nursery/floriculture).

Key Findings

Droughts occurred in the Southern region in 2006, 2008, 2015 and 2019. These events offer insights on the impacts to agriculture and the natural environment, and the potential resilience of communities to future droughts.

Drought resilience was determined by analysing the potential drought impact (risk) and adaptive capacity of each of the twelve LGAs in the Southern region. This showed Clarence had higher adaptive capacity potential to drought, whereas Central Highlands and Derwent Valley display lower adaptive capacity to drought. The potential drought impact (risk) showed Central Highlands with higher risk rating, while Kingborough was lower based on the Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES) Community Vulnerability & Resilience to Drought Index (CVRDI). The potential vulnerability of LGAs in the Southern region to drought impacts are on the lower end of the index.

Therefore, the Southern region has moderate resilience to manage future drought conditions (Figure ES-1). Central Highlands, Glamorgan-Spring Bay and Tasman LGAs have lower resilience to drought and Kingborough, Sorell and Clarence have higher resilience to drought.

However, it is important to note that climate change will increase the frequency, severity and duration of extreme events such as periods of intense heat or rainfall. While the drought risk may be projected to moderately increase in most of the Southern region, it is going to be one of many factors that land managers and communities need to prepare for and respond to in the future.

While agriculture, forestry, and fisheries are not major drivers of the economy in the Southern region in terms of employment, they are important drivers for individual councils (such as Huonville and Southern Midlands), where employment rates in these sectors are high. As the climate continues to change these industries need to continue to adapt and transform to ensure they are resilient to drought and other changes in climate.

Climate change is already impacting on agriculture and the natural environments and communities on which it relies. If communities can increase their adaptive capacity and resilience to future drought events, then it will also assist in increasing their resilience to other extreme events. It is important to note that more frequent, longer duration and severe droughts may reduce adaptive capacity.

Much work has been done to provide secure water supplies for agriculture in parts of the region, through irrigation schemes and individual landowner investments in storage dams. In general, the region also has a good diversity of agricultural enterprises, both at the property and regional levels. These are two important factors that assist with the region's existing resilience to drought. The development of the Regional Drought Resilience Plans will help to identify regional needs, priorities and challenges and inform future investment to improve economic, social and environmental resilience to drought.




Figure ES-1: Drought resilience of Southern region LGAs.

Recommendations

Based on the key findings in this report the following recommendations have been identified.

Table ES-1: Recommendations.

Theme	Recommendation
Use this report to inform community engagement undertaken by the Regional Project Coordinator	<ol style="list-style-type: none"> 1. Test the risk, adaptive capacity and resilience to drought findings by LGA with the community to see if it reflects on-ground experience to past events. 2. Investigate the My Climate View forecast data as a tool to support further community engagement, which explores future climate predictions for individual towns and provides a specific snapshot of how conditions will change in the coming years.
Utilise the data in this report to inform the development of the Regional Drought Resilience Plan	<ol style="list-style-type: none"> 3. Undertake win-win, no regrets actions to assist with short to medium-term adaptation to future drought conditions (i.e. avoid maladaptation). 4. Ensure the Regional Drought Resilience Plan develops long-term transformative actions that consider all five capitals; physical, natural, financial, human and social. 5. Prioritise action in those communities with lower resilience by building adaptive capacity and reducing vulnerability to potential impacts of drought. This includes Central Highlands, Glamorgan-Spring Bay and Tasman LGAs in the Southern region. 6. Continue to develop irrigation schemes where feasible that balance environmental water needs. This is a priority in the Southern Midlands area. 7. Work with land managers to continue to improve and diversify their agricultural operations to be more adaptable to changing climatic conditions as well as extreme climate events (such as drought). This will have broader benefits for agriculture, the natural environment and communities. For example, this may include sustainable agriculture practices that improve soil health through increasing organic matter inputs and reducing losses for greater soil moisture retention. 8. Build technical literacy in regions to enable land managers to utilise current and emerging technology to better plan and prepare for changes in seasonal conditions. 9. Ensure actions consider community health, including mental health, as important aspects of resilience in the Southern region. Be aware that rates of mental health disorders are likely to be higher than are reported.
Undertake monitoring, evaluation, reporting and learning (MERL) for the Regional Drought Resilience Plan	<ol style="list-style-type: none"> 10. Establish clear, measurable and robust indicators of drought resilience in the Regional Drought Resilience Plan, informed by this report and emerging best-practice research. 11. Monitor drought resilience over time and update the Regional Drought Resilience Plan as required, including supporting data.

Glossary

Definition of terms used in the report. Most definitions are sourced from the Intergovernmental Panel on Climate Change¹.

Adaptive Capacity – the combination of the strengths, attributes, and resources available to an individual, community, society, or organisation that can be used to prepare for and undertake actions to reduce adverse impacts, moderate harm, or exploit beneficial opportunities.

Climate Projection – A projection of the response of the climate system to emissions or concentration scenarios of greenhouse gases and aerosols, or radiative forcing scenarios, often based on simulations by climate models. Climate projections are distinguished from climate predictions in order to emphasise that climate projections depend upon the emission/concentration/radiative-forcing scenario used, which are based on assumptions concerning, e.g., future socioeconomic and technological developments that may or may not be realised and are therefore subject to substantial uncertainty.

Drought Resilience – Drought resilience is the ability to adapt, reorganise or transform in response to changing temperature, increasing variability and scarcity of rainfall and changed seasonality of rainfall, for improved economic, environmental and social wellbeing

Impacts – effects on natural and human systems.

Risk – effect of uncertainty on objectives²

Vulnerability – The propensity or predisposition to be adversely affected.

Acronyms

ABARES – Australian Bureau of Agricultural and Resource Economics and Sciences

ABS – Australian Bureau of Statistics

BoM – Bureau of Meteorology

CMIP5 - Coupled Model Intercomparison Project Phase 5 data

CMIP6 - Coupled Model Intercomparison Project Phase 6 data

CSIRO - Commonwealth Scientific and Industrial Research Organisation

EVAO - Estimated Value of Agricultural Operations

IPCC – Intergovernmental Panel on Climate Change

LGA – Local Government Area

MCA – multi-criteria analysis

NRE – Department of Natural Resources and Environment Tasmania

NRM – Natural Resource Management

SEIFA - Socio-Economic Indexes for Areas

SRES – IPCC third assessment report; Special Report on Emissions Scenarios from 2000

¹ IPCC, 2012, Glossary or term, In: *Managing Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC).

² Australian Standard, 2015. ISO 14001:2015 Environmental Management

1. Introduction

1.1. Assessing the impacts of drought

Over the coming decades, climate change is expected to bring increasing temperatures, higher evaporation, changes to rainfall patterns and worsening fire conditions to much of south-eastern Australia. These changes are likely to lead to more frequent, severe and prolonged droughts in many parts of the country. Climate variability and drought are among the many risks faced by agricultural and regional communities. Approaches to building drought resilience in these communities must be tailored not only to local weather conditions and climate forecasts, but also regionally specific socio-economic and cultural factors.

The purpose of this report is to provide a snapshot in time of the indicative and potential drought impacts for the Southern region of Tasmania. It answers the following three questions:

1. What is the prevalence, severity and impacts of drought? (Past/current)
2. What is the likely prevalence, severity and impact of droughts? (Future)
3. What are the vulnerabilities, gaps in preparedness and adaptive capacity for drought and other related permanent transitions to a changing climate? (Analysis)

1.2. Project Purpose

The purpose of this project was to deliver data analysis reports of drought risk, resilience, and adaptive capacity for the three regions of Tasmania – North, North West and South. This Southern region drought report will support the engagement activities of the Regional Project Coordinator and underpin the development of the Regional Drought Resilience Plan for the Southern region with regional stakeholders. This report is designed to facilitate conversations with local communities and sense test whether the data aligns with actual on ground observations, when the Regional Drought Resilience Plan is being developed. The project is being completed under the Tasmanian Government's Regional Drought Resilience Program (RDRP), which is jointly funded under the Australian Government's Future Drought Fund (FDF) and the Tasmanian Government.

Identified long-term outcomes of the Drought Resilience Plans include:

- More primary producers preserve natural capital while also improving productivity and profitability
- Stronger connectedness and greater social capital within communities, contributing to wellbeing and security
- Communities implement transformative activities that improve their resilience to drought.

1.3. Definition of drought

There are no universal definitions of drought, although in its most simple form it can be described as a prolonged, abnormally dry period when the amount of available water is insufficient to meet normal use³. How drought interacts with and affects different regions, agricultural enterprises and communities can vary greatly. Ultimately the way in which water is used determines the types of drought that may be occurring as well as the impact it will have.

Four common categories used to measure drought are defined below and used in this report⁴:

- **Meteorological drought** – this is based on the degree of dryness (often in comparison with some 'normal' or average amount) and the duration of the dry period.
- **Hydrological drought** – this is related to the effects caused by periods of low rainfall and its impact on surface and ground water supply which also affects dam and lake levels.

³ BoM, Understanding Drought. <http://www.bom.gov.au/climate/drought/knowledge-centre/understanding.shtml>, accessed 2 August 2023

⁴ White, D. H. & Walcott, J.J (2009). The role of seasonal indices in monitoring and assessing agricultural and other droughts: a review. CSIRO Publishing, Crop & Pasture Science 60: 599-616

- **Agricultural drought** – this drought classification links both the characteristics of meteorological and hydrological drought to agricultural effect through soil water deficits and plant growth requirements. A plant's demand for water is dependent on prevailing weather conditions, biological plant requirements, its stage of growth and the properties of the soil. Hence an agricultural drought severity could vary within a specific region across different agricultural enterprises, depending on the requirements of the particular crop. Agricultural drought impacts livestock enterprises through detrimental effects on livestock productivity, feed costs and quality and availability of pasture.
- **Socioeconomic drought** – this drought classification associates the supply and demand of economic goods and services with elements of meteorological, hydrological, and agricultural drought. It also includes other factors such as prices of inputs, commodities, and management skill, and these may impact on land managers and the broader community.

Drought resilience is the ability to adapt, reorganise or transform in response to changing temperature, increasing variability and scarcity of rainfall and changed seasonality of rainfall, for improved economic, environmental and social wellbeing.⁵ This report analyses the resilience of agricultural, natural environment and community systems to drought.

1.4. Assessment framework

The assessment framework for this project was based around the three key project questions outlined in Section 1.1 above. Each project question has been considered for three areas: agriculture, natural environment and communities. Table 1-1 outlines the relevant data source for each section of the report.

There was a wide range of data available that informed this report. The first step was identifying the most relevant data to use and how current it was, for example climate scenario data is regularly updated. In Tasmania, Climate Future Scenario modelling was conducted by the Antarctic Climate & Ecosystems Cooperative Research Centre in 2010. This work utilises the A2 (2.0-5.4°C warming) and B1 (1.1-2.9°C) scenarios from the Intergovernmental Panel on Climate Change (IPCC) third assessment report; Special Report on Emissions Scenarios (SRES) from 2000. These Climate Futures Reports, although slightly dated, still provide very relevant snapshots and information on likely future climate change impacts, especially under the more likely A2 scenario based on current greenhouse gas emissions trajectories.

A more recent, and also widely used climate model, is the Coupled Model Intercomparison Project Phase 5 (CMIP5) data. This data was first published in 2013. This data is available via LISTmap and models future climate variables such as mean rainfall, maximum and minimum temperatures, extreme heat days, it has also been used for their future enterprise suitability mapping. The CMIP5 data has also been used at a national level for the My Climate View data developed by the Commonwealth Scientific and Industrial Research Organisation (CSIRO) and Bureau of Meteorology (BoM).

It is noted that the above two sets of data have been used interchangeably in this report, based on which source provided the most comprehensive and workable data to use for the specific topic.

The Tasmanian Government will be developing new fine-scale or “downscaled” climate projections for Tasmania, using the latest global climate models (CMIP6). The updated projections will be available in mid-2025 and will support business, industry, community and government to understand the projected future climate for Tasmania, and plan for and build resilience to the impacts of a variable and changing climate.

Other data sources used in this report include the Tasmanian Governments Profile ID dashboard and Australian Bureau of Statistics (ABS) Census data.

⁵ Department of Agriculture, Water and the Environment (2020) Future Drought Fund (Drought Resilience Funding Plan 2020 to 2024) Determination 2020, https://ehq-production-australia.s3.ap-southeast-2.amazonaws.com/a69d99c1b753c9e93cdf88ce1fd6d723997f78d9/original/1584927882/Drought_Resilience_Funding_Plan.pdf_7c8e152b3f40ab97f9e17b39fc4ce42a?X-Amz-Algorithm=AWS4-HMAC-SHA256&X-Amz-Credential=AKIA4KKNQAKIOR7VAOP4%2F20231016%2Fap-southeast-2%2Fs3%2Faws4_request&X-Amz-Date=20231016T023846Z&X-Amz-Expires=300&X-Amz-SignedHeaders=host&X-Amz-Signature=20de2119787c15ed7867f67c210e8f670c91122beb19b9d4522b7c3014993942

Table 1-1: Data sources used in report, by report section.

Section	Data Source	Data Access
2.1	Profile ID Tasmania, utilising ABS Census data	https://profile.id.com.au/tasmania
	ABS Estimated Value of Agricultural Operations through the Agricultural Commodity Survey	https://www.abs.gov.au/statistics/industry/agriculture/value-agricultural-commodities-produced-australia/2021-22
2.2	Natural Resource Management (NRM) South Strategy to 2030	https://nrmsouth.org.au
	LISTmap by the Department of Natural Resources and Environment Tasmania (NRE)	https://maps.thelist.tas.gov.au/listmap/app/list/map
2.3	Profile ID Tasmania, utilising ABS Census data	https://profile.id.com.au/tasmania
3	Climate Change Australia	https://www.climatechangeinaustralia.gov.au/en/changing-climate/state-climate-statements/tasmania/
	Bureau of Meteorology historical rainfall maps for Tasmania	http://www.bom.gov.au/climate/maps/rainfall/
	NRE Annual River Reports	https://nre.tas.gov.au/water/water-data/annual-river-reports
3.1	The Long Paddock, Queensland Government	https://www.longpaddock.qld.gov.au/
3.2-3.4	Bureau of Meteorology – Previous Droughts	http://www.bom.gov.au/climate/drought/knowledge-centre/previous-droughts.shtml
	ABC Rural – Various Articles	https://www.abc.net.au/news/rural
4.1	Climate Change Australia	https://www.climatechangeinaustralia.gov.au/en/changing-climate/state-climate-statements/tasmania/
	The Antarctic Climate & Ecosystems Cooperative Research Centre – Technical Reports	https://climatefutures.org.au/projects/climate-futures-tasmania/
4.2	My Climate View	https://myclimateview.com.au/
	LISTmap by NRE – Mean rainfall	https://maps.thelist.tas.gov.au/listmap/app/list/map

Section	Data Source	Data Access
	The Antarctic Climate & Ecosystems Cooperative Research Centre – Technical Reports	https://climatefutures.org.au/projects/climate-futures-tasmania/
4.3	Climate Change Australia	https://www.climatechangeinaustralia.gov.au/en/projections-tools/climate-analogues/
	LISTmap by NRE – Enterprise Suitability Mapping	https://maps.thelist.tas.gov.au/listmap/app/list/map
4.4	NRM South Strategy to 2030	https://nrmsouth.org.au
4.5	The Antarctic Climate & Ecosystems Cooperative Research Centre – Technical Reports	https://climatefutures.org.au/projects/climate-futures-tasmania/
	Profile ID Tasmania, utilising ABS Census data (employment)	https://profile.id.com.au/tasmania
5.1	Socio-Economic Indexes for Areas (SEIFA)	https://www.abs.gov.au/websitedbs/censushome.nsf/home/seifa
	ABS Estimated Value of Agricultural Operations through the Agricultural Commodity Survey	https://www.abs.gov.au/statistics/industry/agriculture/value-agricultural-commodities-produced-australia/2021-22
	Profile ID Tasmania, utilising ABS Census data	https://profile.id.com.au/tasmania
	Tasmania Irrigation – Existing Irrigation Schemes	https://www.tasmanianirrigation.com.au/active-schemes-map
	LISTmap by NRE – Existing Dams	https://maps.thelist.tas.gov.au/listmap/app/list/map
5.2	ABARE’s Community Vulnerability & Resilience to Drought Index	https://www.agriculture.gov.au/abares/research-topics/climate/drought/resilience#community-vulnerability-and-resilience-to-drought-index-cvrdi_2
5.3	Data from Sections 5.2 and 5.3 combined	
5.4	RMCG & Australian Resilience Centre (2020) Goulburn Murray Resilience Strategy	https://www.rdv.vic.gov.au/resources/resilience

1.5. Quantitative analysis

As part of the project, quantitative data has been utilised to develop a multi-criteria analysis (MCA) that maps the indicative regional drought impact (risk), adaptive capacity and resilience. The framework used to inform this assessment is the ABARES adopted Schematic of Drought Sensitivity, Risk and Resilience Model (see Figure 1-1).




Figure 1-1: Drought Risk, Adaptive Capacity and Resilience Model⁶.

The drought impact (risk) data was sourced from the ABARES Community Vulnerability and Resilience to Drought Index (CVRDI) which has calculated the potential drought impact for each LGA in Australia. To determine the Adaptive Capacity of each LGA in the region, at least two data sources were identified for the five capitals (physical, natural, financial, human and social) and were then ranked. The drought impact score and the adaptive capacity score for each LGA were then combined to identify an indicative measure of drought vulnerability and resilience for each LGA within the region. The results of the assessment are presented in Section 5, and the full assessment methodology is provided in Appendix 1.

In addition to the five capitals in the adaptive capacity box in Figure 1-1, is Economic Diversity. This refers to the spread of employment across different industries. This has not been included in this version of the MCA as an individual indicator for adaptive capacity, as it was determined that employment information was adequately captured in the community sensitivity analysis as well as in financial and human capital. However, if deemed appropriate it could be included in a future version of the MCA.

⁶ <https://www.agriculture.gov.au/abares/research-topics/climate/drought/resilience#reports>

2. Regional Profile

2.1. Overview

The Southern region has a land area of 25,964 km², a population of just under 300,000 and is comprised of the following Local Government Areas (LGAs):

- Brighton
- Central Highlands
- Clarence
- Derwent Valley
- Glamorgan-Spring Bay
- Glenorchy
- Hobart
- Huon Valley
- Kingborough
- Sorell
- Southern Midlands
- Tasman.




Figure 2-1: The Southern region and LGAs.

The data analysed in this section has been drawn from ABS 2021 Census data and manipulations of this data available through Profile ID. Data has been selected to give a brief overview of the overall demographics of the area and also assist with understanding the possible resilience vulnerabilities and strengths that may exist within the population of the Southern region.

Compared to Tasmania as a whole, households in the Southern region are skewed towards the higher end of household incomes, people tend to be more educated and slightly younger. Rent and mortgage repayments are higher than Tasmania as a whole. A smaller proportion of people own their own home, while rental rates are about the same. People are more likely to work in education and training, public administration & safety and professional, scientific and technical services and less likely to work in retail trade, manufacturing and primary industries.

Clarence, Hobart and Kingborough have the least level of disadvantage of the LGAs in the Southern region, while people within the Brighton and Central Highlands LGA demonstrate the most disadvantage.

Only 3.5% of the population in the Southern region are employed in agriculture, forestry and fisheries. However, this increases in some LGAs, such as Huon Valley and the Southern Midlands where these sectors employ around a quarter of the population. Unsurprisingly Huon Valley and Southern Midlands also account for around 41% of the region's agricultural output. Fruit (including Nuts) is the region's largest contributor to the value of agriculture, this is followed by livestock products (excluding dairy), and Field Crops (including Nursery/Floriculture).

2.2. Agriculture and Industry

2.2.1. Regional Agricultural Production

The primary production system includes agriculture, fisheries and forestry, as well as the infrastructure and workforce supporting and servicing these industries. The primary production industries contribute significantly to state and regional economies, and are major employers and exporters. The agriculture industry includes growers, farmers of livestock, and nurseries. The range of agricultural products and the value of production are highly variable across the landscape.

Agricultural production systems, landowners' choices about the mix of commodities they produce, and their associated profitability are determined by a range of factors, including:

- Natural assets of their land (land capability, climate, access to irrigation water, soil type, slope, drainage, rainfall, temperatures)
- Property shape, geographic location and size
- Personal or business farming experience, preferences or tradition
- Infrastructure and machinery (existing or legacy; costs to purchase, construct and maintain; changing technology; and market/processor requirements)
- Market proximity/accessibility (i.e., processor/packer or purchaser location in relation to property, and ability to transport the commodity that distance)
- Market demands and conditions (seasonal or long term trend), or existing contract requirements
- Ownership or lease structure.

Many farmers operate multiple agricultural enterprises concurrently or in rotation on their land, producing a range of commodities. They rotate the enterprises between paddocks and over time to match market requirements, supply quotas, and for other management reasons, including weed, disease, and pest control.

The primary production sector (agriculture, fisheries and forestry) is a strong contributor to the economy for some LGAs in the Southern region, both in terms of employment (as described above), but also in the value of production.

According to Informed Decisions (ID), in the 2021/22 financial year, the primary production sector added 8% of the overall economic value of the Southern region⁷. In the Southern Midlands, primary production contributed 55% of the LGA's economic value, while in the Huon Valley, it contributed 43%.

The Estimated Value of Agricultural Operations (EVAO) is a measure of agricultural value used by both the Australian Bureau of Statistics (ABS) and the Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES), part of the Commonwealth Department of Agriculture, Fisheries and Forestry (DAFF). The ABS and ABARES are usually the major sources of agricultural and commodity value and distribution in Australia.

The data presented here is based on ABS 2020/21 data. The ABS collected this data through the Rural Environment and Agricultural Commodity Survey. To be included in the survey businesses must have:

- An Australian Business Number (ABN)
- Undertaken agricultural activity
- An EVAO of \$40,000 or greater.

Table 2-1 identifies the value of agriculture of each LGA in the Southern region. This shows that Huon Valley accounts 25.71% of the region's output, with Southern Midlands producing 16.83% and Derwent Valley producing 15.26%. It is important to note that these figures do not include the value of plantation forestry or aquaculture. Table 2-2 shows that fruit (including nuts) is the region's largest contributor to the value of agriculture (9.95%), this is followed by livestock products (excluding dairy) 31.93%, and field crops (including nursery/floriculture) (14.99%).

Table 2-1: Value of agriculture per LGA.

South Region LGA	\$ (million)	Percentage of Region
Brighton	2.18	0.69%
Central Highlands	38.875	12.31%
Clarence	16.204	5.13%
Derwent Valley	48.213	15.26%
Glamorgan-Spring Bay	25.989	8.23%
Glenorchy	0.161	0.05%
Hobart	0.005	0.00%
Huon Valley	81.228	25.71%
Kingborough	8.856	2.80%
Sorell	27.799	8.80%
Southern Midlands	53.152	16.83%
Tasman	13.216	4.18%
Total	315.878	

⁷ <https://economy.id.com.au/tasmania/value-add-by-industry?WebID=400>

Table 2-2: Value of agricultural enterprises in the Southern region.

Agricultural Enterprise Type	\$ (million)	Percentage of Region
Livestock (inc meat products, wool & eggs)	100.86	31.93%
Vegetables	13.83	4.38%
Dairy	13.53	4.28%
Fruit (inc nuts)	126.21	39.95%
Field Crops (inc nursery/floriculture)	47.37	14.99%
Wine	14.08	4.46%
Total	315.88	

2.2.2. Employment

The 10 industry sectors in Table 2-3 account for 80% of the total employed persons over the age of 15 in the Southern region. Sectors showing large differences to Tasmania as a whole are highlighted, with blue indicating a greater proportion of Southern region residents employed in the sector compared to the whole state and tan indicating a lesser proportion. Of the other 10 sectors there was proportionately little difference between the Southern region and Tasmania as a whole with the exceptions of: a greater proportion in the Arts and Recreation Services sector, and relatively fewer in the Transport, Postal & Warehousing, Wholesale Trade and Mining sectors. However, when considering different Council areas in the Southern region, Table 2-4 the percentage of the workforce that work in the agriculture sector varies greatly across the Councils. Table 2-5 shows approximate number of people employed in each sub-category of agriculture. It is noted the Aquaculture is a major employer across a number of LGAs.

Table 2-3: Major employment sectors.

Industry Sector	People	Southern Region	Tasmania
Health Care and Social Assistance	22,091	16.2%	16.4%
Education and Training	13,890	10.2%	9.4%
Public Administration and Safety	12,943	9.5%	7.3%
Retail Trade	12,351	9.1%	9.6%
Construction	11,806	8.7%	8.6%
Accommodation and Food Services	10,689	7.8%	7.6%
Professional, Scientific and Technical Services	8,437	6.2%	5.1%
Manufacturing	7,134	5.2%	6.4%
Other Services	4,971	3.7%	3.8%
Agriculture, Forestry and Fishing	4,822	3.5%	5.3%

Table 2-4: Percentage of the workforce employed in the Agriculture Sector by LGA, compared to regional and state percentages (2021).

LGA	Agriculture, Forestry and Fishing
Brighton	3%
Central Highlands	8%
Clarence	3%
Derwent Valley	12%
Glamorgan-Spring Bay	18%
Glenorchy	1%
Hobart	1%
Huon Valley	24%
Kingborough	5%
Sorell	8%
Southern Midlands	29%
Tasman	18%
Southern Tasmania (region)	3.5%
Tasmania (state)	5.3%

Table 2-5: Number of people employed in the agriculture industry by subsector.

LGA	Agriculture	Aquaculture	Forestry and Logging	Fishing, Hunting and Trapping	Agriculture, Forestry and Fishing Support Services
Brighton	111	0	5	0	19
Central Highlands	315	30	12	0	16
Clarence	359	141	8	78	22
Derwent Valley	233	24	97	0	39
Glamorgan-Spring Bay	171	136	24	41	18
Glenorchy	71	55	34	23	14
Hobart	126	353	45	64	44
Huon Valley	551	803	49	54	41
Kingborough	182	297	4	67	15
Sorell	226	62	8	21	8
Southern Midlands	380	0	0	0	20
Tasman	97	50	9	24	0
	2822	1951	295	372	256

Additional economic data on income, housing and education is provided in Appendix 2.

2.3. Natural Environment

2.3.1. Bioregions

'Bioregions are large, geographically distinct areas of land with common characteristics such as geology, landform patterns, climate, ecological features and plant and animal communities.'⁸

There are nine bioregions in Tasmania. The South East and Southern Ranges bioregions dominate the Southern region. The Tasmanian West covers the western parts of the Derwent Valley and Huon Valley LGAs. The Central Highlands bioregion covers the northern half of the LGA of the same name. And the southern tip of the Northern Midlands bioregion extends into the Southern Midlands LGA.

⁸ <https://www.dcceew.gov.au/environment/land/nrs/science/ibra/australias-bioregion-framework>

2.3.2. Flora and Fauna

TASVEG 4.0 lists 11 vegetation community groups. In the Southern region, modified land occurs almost continuously down the east coast except on conserved or hillier areas where there is dry eucalypt forest and woodland interspersed with native grassland with occasional areas of scrub, heathland and coastal complexes. The eucalypt forest grades to wet on higher ground. Large swathes of modified land also occur along the south east coast and stretch inland towards the Midlands along the major river valleys, and continue south of Hobart to the mouth of the Huon River. West and south of Huonville wet eucalypt forest and woodland tend to predominate. This grades to communities of wetter and/or higher regions such as moorland, sedgeland and rushland communities and highland and treeless vegetation towards the higher and/or wetter regions of the Central Plateau / Tasmanian Wilderness World Heritage Area in the west and south west. Notably a long stretch of modified land just to the east of the World Heritage Area is attributable to silviculture rather than agriculture. In the north west of the region there are large areas of native grassland in amongst dry eucalypt forest and modified land, these become smaller towards the Midlands and the south.

In the Southern region, there are large areas of modified land, much of it on the low lying plains of the region, including down the east and south east coasts.

There are several threatened ecological communities within the region.⁹ *Eucalyptus ovata* forest is listed as vulnerable. It is found in the northern region of the Glamorgan-Spring Bay LGA¹⁰. The endangered Alpine sphagnum bogs and associated fens are found in elevated areas throughout the region, particularly around the Central Plateau, while the similarly endangered giant kelp marine forests of south east Australia community is found around most of the Southern region coast, including the major offshore islands.

Lowland native grasslands are listed as critically endangered. This ecological community is particularly prevalent in the Northern Midlands LGA and has contributed to the recognition of the Midlands biodiversity hotspot as one of only 15 in Australia. It is less prevalent in the Southern Midlands LGA. It is present in most of the LGAs of the Southern region, particularly Brighton. It is not recorded in the Huon Valley or Glenorchy LGAs.

There are three further ecological communities. Subtropical and temperate coastal saltmarsh (vulnerable) occurring in patches along the extent of the east coast to Recherche Bay; Tasmanian forest and woodlands dominated by black gum or Brookers gum (*E. ovata/E. brookeriana*) and Tasmanian white gum (*E. viminalis*) wet forest are both either found or likely to occur across the region. The first of these is listed as vulnerable while the last two are critically endangered ecological communities.

Within the Southern region there are 261 species that have 100% of their recorded range within the region. There are a further 304 species with greater than 50% of their recorded range within the region.¹¹ 114 species occurring in Southern region are listed as threatened.¹²

2.3.3. Tasmanian Wilderness World Heritage Area

Much of the western edge of the Southern region is in the Tasmanian Wilderness World Heritage Area and spans large portions of the Huon Valley, Derwent Valley and the Central Highlands LGAs.

2.3.4. Catchments

At its essence, the geographic area where rain falls and drains to is a catchment, however the boundaries between these can be hard to determine. There are 48 major catchments within Tasmania¹³, 13 are wholly

⁹ <https://www.dcceew.gov.au/environment/biodiversity/threatened/communities/tas>

¹⁰ From DCCEEW's Protected Matters search tool - <https://www.dcceew.gov.au/environment/epbc/protected-matters-search-tool>

¹¹ <https://www.dcceew.gov.au/parks-heritage/heritage/publications/anhat/biodiveristy-summary/south-tas>

¹² <https://nrmsouth.org.au/wp-content/uploads/2023/04/NRM-South-Regional-Strategy-2030.pdf>

¹³ <https://nre.tas.gov.au/water/a-guide-to-water-in-tasmania/tasmanias-water-catchmentstasmanian>

or substantially within the Southern region and four are partially within the region. The largest catchment is the Gordon-Franklin, that spans the Southern and North West regions, and the smallest is Great Lake.

2.3.5. Wetlands

There are 65 internationally listed (Ramsar) wetlands in Australia, 10 are in Tasmania with four in the Southern region – Moulting Lagoon and the nearby Apsley Marshes, Pittwater-Orielton Lagoon, and Interlaken. Tasmania also has 89 wetlands recognised for their national importance and listed in the Directory of Important Wetlands in Australia¹⁴. The wetlands, which can be permanent, seasonal or ephemeral, and vary from one to over 27,000 hectares in size. They are found in a wide range of locations, including alpine, moorland, freshwater, estuarine and subterranean environments.

Within the Southern region wetlands of very high conservation value are scattered across the region, with large areas in the Central Highlands LGA, and throughout the World Heritage Area, and areas around the Apsley Marshes, the Derwent and Huon Rivers, Mount Wellington, Bruny Island, and the Tasman and Forestier Peninsulas. There are further smaller wetlands of very high conservation within the region. There are many wetlands of all classifications in the Southern region however there are relatively fewer in areas more likely to be subject to agriculture or urbanisation within the region.

2.3.6. Climate

The Southern region is temperate with mild/warm summers and cold winters. Unlike the rest of the Tasmania, the Southern region has uniform rainfall (500-800 mm/yr) across the year (compared to a marked wet winter and dry summer). There are small areas in the south west with cool summers and no dry season.

To the west (and to the north over the Central Highlands) there is more rainfall (>800 mm rainfall/yr) and a gradation to a marked wet winter.

2.3.7. Land Capability

The ability of land to support agriculture is divided into seven classes of land capability. Classes 1 to 3 are considered prime agricultural land in Tasmania, 4 is considered marginal for cropping and Class 7 is the poorest land.¹⁵ There are only a few small pockets of land in the Southern region that are mapped as Class 3 land, these are located at Marion Bay, Campania-Runnymede, Kingston and Grove. The balance of agricultural land is generally a mix of Class 4, Class 5 and Class 6 land.

2.4. Community

Census data allows for three measures of population:

- Enumerated (who was in the area on the night)
- Usual residence (where a person usually lives)
- Estimated resident population (calculated after the Census to take into account those who missed the Census or were overseas at the time; updated annually).

The estimated resident population for the Southern region (2022): 296,173

The usual resident population for the Southern region (2021): 288,818

The Southern region has a lower median age than Tasmania (40 compared to 42).

¹⁴ <https://nre.tas.gov.au/conservation/flora-of-tasmania/tasmanias-wetlands>

¹⁵ <https://nre.tas.gov.au/agriculture/land-management-and-soils/land-and-soil-resource-assessment/land-capability/the-land-capability-classification-system>

Table 2-6: Key population statistics for the Southern region.

Key population statistics	Southern Region	Tasmania
Population density (people / km ²)	11.4	8.41
Median age	40	42
Aboriginal and Torres Strait Islander population	5.0%	5.4%
Religious belief	43.4%	43.4%
People needing assistance with core activities	6.3%	6.8%

Religious belief and those requiring assistance with core activities have been included in Table 2-6 as they may contribute to personal resilience strengths and vulnerabilities. More people under 19 and over 70 are requiring assistance than at the time of the previous census (2016). There are over 18,000 people that need assistance with core activities, a rise of over 2,400 people from the previous census.

Table 2-7: Country of birth and languages other than English (LOTE) in the Southern region.

Key population statistics	Southern Region	Tasmania
Overseas born	18.0%	15.4%
LOTE at home	11.5%	8.7%
Speaks English not well or not at all	3,166 (1.1%)	0.9%
Speaks English well or very well	30,567 (10.6%)	8.0%

As communications with key stakeholders may be a consideration in Drought Resilience Planning, the following information concerning country of birth and languages spoken at home is included.

There is a significant cohort of German, Dutch and Italian born people within the Southern region (10,801, 5,728 and 5,092 people respectively). However, of these languages, only German is listed as being spoken at home by a large number of people (less than 1,000 individuals or 0.3% of the population).

The Chinese-born population (9,331 people in 2021, 3.2% of the Southern region population) is the largest group born in a country where the first language is unlikely to be English. Almost 6,500 people speak Mandarin at home (6,482) and around 1,200 speak Cantonese (1,198).

There are slightly more people speaking Nepali at home (4,710) than those born there (4,271). Of the Indian born population (4,179), around half speak Punjabi at home, and a quarter each Hindi or Urdu. Those born in Vietnam and the Philippines account for about a 1,000 people each, with about 1,000 people speaking Vietnamese at home, and 800 Filipino/Tagalog.

The Wicking Dementia Research and Education Centre within the University of Tasmania has been conducting the Island Study Linking Ageing and Neurodegenerative Disease (ISLAND) project since 2019. This longitudinal study collects data from those over 50 who live in Tasmania. Participants are asked a wide range of questions including those relating to the Lubben Social Network Scale which measures the perceived social support of family and friends. Statewide, the Wicking Centre has around 1,000 people willing to share their data outside the ISLAND project. All participants are over 50 years of age (median and mean both approximately 66) and mostly female (around 74%).

Residents of Derwent Valley LGA self-report as the most socially isolated within the southern region, while Hobart and Kingborough residents report as the most socially supported. Clarence, Kingborough and Tasman LGAs also report relatively good social connectivity.

Respondents represent between 0.32% and 1.38% of the population of their LGA. Brighton, Glenorchy, Sorell and Huon Valley LGAs fall between these bounds.

Glamorgan-Spring Bay reported the highest levels of social support but from only 19 participants (0.37% of the LGA's population). Similarly, Southern Midlands residents report feeling low social support, however this is from numbers lower than those for Glamorgan-Spring Bay. Central Highlands had too few respondents to include.

3. Past and Current Impacts of Drought

Tasmania has been experiencing impacts from climate change¹⁶ over the past 100 plus years that have impacted the frequency of droughts. This includes:

- Average annual temperature has increased by 1.1°C since 1910
- Decrease in annual rainfall since 1900
- Increase in the number of dangerous bushfire days where wind, humidity, temperature and precipitation levels favour fire.

Bureau of Meteorology historical rainfall maps for Tasmania (drought) from 2000 to 2022 (Figure 3-1) indicates that there have been a number of low rainfall years where drought has occurred in this timeframe. For areas of the Southern region the most prominent drought calendar years have been 2006, 2008, 2015 and 2019

¹⁶ CSIRO, <https://www.climatechangeinaustralia.gov.au/en/changing-climate/state-climate-statements/tasmania/>, accessed 9/8/23




Figure 3-1: Annual low rainfall years 2000-2021 (source: BoM).

3.1. Past Climate Trends for Key Townships

This section provides climate details on four representative townships in the Southern region: Bothwell, Huonville, Richmond and Triabunna. The climate details provided include data from 1991 to 2020 on temperature, rainfall and evapotranspiration. Rainfall deficit data compares annual rainfall data and annual evapotranspiration and is a good indicator of annual soil moisture levels. The data generally shows that the BoM annual drought data aligns with the township data.







Figure 3-2: Annual temperatures (°C) from 1991 to 2020 (source: Long Paddock).







Figure 3-3: Annual rainfall (mm) from 1991 to 2020 (source: Long Paddock).







Figure 3-4: Annual evapotranspiration (Eto) (mm) from 1991 to 2020 (source: Long Paddock).







Figure 3-5: Annual deficit – rainfall minus evapotranspiration (mm) (source: Long Paddock).

3.2. Impacts on Agriculture

In general, the agricultural areas of the Southern region have relatively low annual rainfall levels compared to the rest of Tasmania. So, in lower-than-average rainfall years the impact on agricultural productivity is high. This has impacts on feed availability for livestock, and can mean land managers are required to destock, and it will also impact on yields of both perennial and annual crops. In 2018 it was reported that the East Coast was in significant need of assistance for feedstock, due to dry conditions and rising feed costs. This came after many of the farmers who were seeking aid had in previous years sent feed to parts of mainland Australia to provide relief to farmers in drought-stricken areas.

From September 2007 to March 2009 the Federal Government formally declared that areas of northern Tasmania required Exceptional Circumstances (EC) drought assistance¹⁷. The areas identified were:

- The northern and western portions of the Southern Midlands LGA
- The Coal River Valley in the Clarence and Sorell LGA
- The north west portion of the Brighton LGA and a small northern portion of the Derwent Valley LGA
- The eastern Central Highlands as far west as Wayatinah
- A buffer zone along the East Coast extending from St Marys (Northern region) to Primrose Sands (Southern region).

In 2016 the Tasmanian and Federal Government announced a Drought Concessional Loan Scheme for Tasmanian farming businesses. Eligible businesses were able to apply for a loan of up to \$1 million to be used for restructuring debt, to fund operating expenses, or drought preparedness and recovery activities¹⁸.

In some areas of the Southern region, land managers are experiencing greater water security as more irrigation scheme water comes online, assisting with offsetting some of the impacts of drought on agriculture.

Combined pressure from climate variability and commodity prices can cause significant financial stress for agricultural businesses. In simple terms, research has demonstrated that the most profitable years for farmers tend to be high rainfall years with favourable commodity prices, and the least profitable tend to be drought years with unfavourable prices¹⁹. This relationship can be complex however, as high commodity prices due to drought-induced scarcity can benefit farmers who experience lower impacts and are still able to produce, creating a polarised 'winners and losers' situation²⁰. This divergence tends to become more exacerbated as droughts increase in severity and duration.

At the farm level, the impacts of drought can be complex, involving changes to productivity, outputs and input prices and their subsequent interactions. Changes to labour requirements, fodder transport, machinery efficiency, input use, pumping and irrigation requirements all affect how individual farmers fare in drought years²⁰. Drought impacts can extend beyond the geographical limits of low-rainfall weather conditions through influencing markets. For example, livestock farmers may consider themselves drought affected even when experiencing average local weather conditions, if drought in other regions leads to higher feed prices²¹.

¹⁷

<https://parlinfo.aph.gov.au/parlInfo/search/display/display.w3p;orderBy=customrank;page=0;query=drought%20declaration%20tasmania;rec=2;resCount=Default>

¹⁸

https://www.premier.tas.gov.au/site_resources_2015/additional_releases/drought_concessional_loans_now_open_for_tasmanian_farm_businesses

¹⁹ Hughes, N, Galeano, D and Hatfield-Dodds, S 2019, *Analysis of the effects of drought and climate variability on Australian Farms*, ABARES Insights Issue 6, Department of Agriculture, Australian Government.

²⁰ Fleming-Muñoz, D.A., Whitten, S. and Bonnett, G.D 2023, *The economics of drought: A review of impacts and costs*, Australian Journal of Agricultural and Resource Economics.

²¹ Hughes, N, Soh, WY, Boulton, C and Lawson, K 2022, *Defining drought from the perspective of Australian farmers*, Climate Risk Management, Volume 35.

Drought is also experienced differently across agricultural industries such as cropping or livestock production¹⁹. Livestock enterprises can experience drought effects after the actual drought period when restocking and herd rebuilding require significant expenses²². Cropping enterprises may face a greater risk from climate variability than livestock, with negative impacts driven by lower crop yields and less area planted during drought periods¹⁹. However, cropping may rebound more quickly when rainfall increases following drought²⁰.

3.3. Impacts on the Natural Environment

In times of drought, water scarcity affects entire ecosystems. During times of prolonged lack of rain, there are interdependent landscape-wide effects. Decreased soil moisture affects all vegetation, including natives and non-irrigated crops and grasses. Native vegetation can become drier than usual, creating greater potential for bushfires. With less available food and water, wildlife will roam further and potentially have a greater impact on agricultural crops. With less groundcover, and projected larger rain events, there is also greater risk of erosion and loss of valuable topsoil.

An example of how extreme heat and dry conditions can affect native flora is 'ginger tree syndrome' which affects some *Eucalyptus* species such as *Eucalyptus viminalis* (white gum). Over the 2013/2014 summer there were widespread reports of this occurring in areas of northern Tasmania. Ginger tree syndrome occurs where prolonged high temperatures and dry conditions cause water stress and heat shrinkage of the bark and trunk of *Eucalyptus* trees²³. The red-coloured sap seeps through the bark, prominently staining affected trees a 'ginger' colour. The syndrome typically causes tree mortality within 12 months and was the cause of widespread tree deaths following the 2013/2014 summer. Ginger tree syndrome is becoming a perennial threat to *Eucalyptus viminalis* across Tasmania. *Eucalyptus viminalis* wet forest is listed as critically endangered under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act).

Drought has many compounding impacts on waterways. Seasonal stream and river flows decrease in drought, reducing the amount of habitat available to aquatic species²⁴. Where flows are reduced, water temperature may rise and lead to a drop in dissolved oxygen, negatively impacting water quality. The adverse impacts of sediments, nutrients and pollutants can also be elevated where flows are reduced. Stream flows are necessary for maintaining riparian vegetation, floodplain and wetland condition and productivity. Under drought conditions, the reduced flows can lead to changes in riparian and floodplain vegetation, which can increase erosion and accelerate land degradation²⁴.

Soil is vulnerable to drought, particularly where land use change and poor land management have already led to reduced soil function and quality. Significant parts of Australia's agricultural landscape already face pressure from erosion, salinisation, soil carbon loss, acidification, contamination and urban expansion, which compound and exacerbate the impacts from drought²⁵. Loss of topsoil, nutrient imbalances, compaction, reduced soil organic carbon and biological activity can place soils at increased risk of erosion and loss of productivity when drought conditions arise²⁴. Soils managed under regenerative strategies that maintain groundcover and minimise compaction and disturbance can be more resilient to drought²⁴.

Increases to average temperatures, changes in precipitation and occurrences of extreme events have the potential to worsen invasive species and disease problems²³.

²² Hooper, S, Ashton, D, Crooks, S, Mackinnon, D, Nicols, P, & Phillips, P 2008, *Farm Financial Performance: Australian Farm Income and Drought Recovery, 2005-06, 2006-07 and 2007-08*, Australian Commodities: Forecasts and Issues.

²³ Department of Agriculture, Water and the Environment 2021, *Conservation Advice for the Tasmanian white gum (Eucalyptus viminalis) wet forest*, Australian Government.

²⁴ Department of Primary Industries, Parks, Water and Environment Resource Management and Conservation Division 2010, *Vulnerability of Tasmania's Natural Environment to Climate Change: An Overview*, Tasmanian State Government.

²⁵ Williams, KJ, Hunter, B, Schmidt, RK, Woodward, E and Cresswell, ID 2021, Land: Soil. In: *Australia State of the Environment 2021*, Department of Agriculture, Water and the Environment, Australian Government.

Extreme events such as droughts, fires and floods can favour adaptable invasive species and lead to sudden increases in weed and pest extent and impact²⁶. Changes in climate are also expected to favour the spread of disease vectors such as mosquitoes, ticks and rodents which have the potential to transmit diseases among wildlife, livestock and humans²³. Wildlife diseases have been emerging in Tasmania at an increased rate since the 1990s, reflecting a national and global trend²³. The risk of increased weed and insect pests and diseases strongly extends to agricultural systems²⁷.

3.4. Impacts on Communities

As with the natural environment, communities also have interdependencies - what happens in one section of the community will have effects on people in other sections of the community.

Droughts, as with other extreme environmental events, cause significant stress on communities. Effects can include health, particularly mental health, and economic/financial impacts. Long standing drought will impact the financial viability of individual farming enterprises, affecting on-farm and regional employment, the available money within the community and the viability of all the businesses that rely on that local expenditure. It will also impact on residential water supplies.

Research has demonstrated that the far-reaching impacts of drought on communities negatively affect mental health²⁸. Disruptions to social and economic systems such as crop and livestock failure, social isolation from increased workload and reduced resources, financial hardship, lack of water and migration out of rural communities can increase stress and trigger mental disorders such as depression²⁸. Farming communities also show resilience to drought through positive coping mechanisms, and evidence suggests that reliance shared social values and resources increases in times of drought²⁹.

At the community scale, drought has significant economic and social costs that play out over the short and longer term³⁰. Substantially lower employment, yet increased farm workloads for all family members, has been recorded. Businesses in rural towns suffer due to local populations earning and spending less and fewer visitors, again leading to employment disruptions. Children and young people also face increased farm labour pressure, which can cause them to miss school and other education opportunities³⁰. A loss of a wide range of services, including banks, schools and medical services, has also been associated with drought³¹. Uncertainty and concern for the future, coupled with the lack of support services, can lead young people to lose interest in farming, resulting in a loss of potential future farmers in communities³⁰.

Members of a community may experience drought differently depending on demographic factors. Aboriginal and Torres Strait Islander Australians are disproportionately impacted by drought, due to persistent health and social disadvantages³⁰. Women report many changes to their roles, taking on additional farm or supplementary work, further care of children and their education, and being the emotional support in their family while placing their own health and wellbeing as a low priority³².

²⁶ Invasive Species Council 2009, *Invasive species and climate change*, accessed 10/10/2023 from <https://invasives.org.au/wp-content/uploads/2014/02/fs_invasivespeciesclimatechange.pdf>

²⁷ International Plant Protection Convention Secretariat 2021, *Scientific Review of the impact of climate change on plant pests – A global challenge to prevent and mitigate plant pest risks in agriculture, forestry and ecosystems*, Food and Agriculture Organisation of the United Nations.

²⁸ Luong, TT, Handley, T, Austin, EK, Kiem, AS, Rich, JL and Kelly, B 2021, *New Insights Into the Relationship Between Drought and Mental Health Emerging from the Australian Rural Mental Health Study*, *Frontiers in Psychiatry* volume 12.

²⁹ Caldwell, K and Boyd, CP 2009, *Coping and resilience in farming families affected by drought*, *Rural and Remote Health* volume 9.

³⁰ Lester, L, Flatau, P and Kyron, M 2022, *Understanding the Social Impacts of Drought*, Centre for Social Impact, University of Western Australia.

³¹ Edwards, B, Gray, M and Hunter, B 2008, *Social impacts associated with drought*, Australian Institute of Family Studies, Australian Government.

³² Alston, M and Kent, J 2004, *Social impacts of drought: a report to NSW Agriculture*, Centre for Rural Social Research, Charles Sturt University.

For young people, a ‘missed childhood’ can be a severe outcome of drought, where they are forced to take on adult roles, increased workloads, and a higher risk of physical injury³³.

Research indicates that men in farming families are less likely to seek help for mental health issues due to stigma and are more likely to withdraw from their communities³⁰. Older members of farming communities suffer from increased workloads and declining physical ability, loss of identity and resistance to using mental health services³⁰.

Tasmania’s electricity is predominantly sourced from hydropower. Long periods of low rainfall impact the water levels of the state’s hydro-electric dams, with flow-on effects to the state’s power generation capacity. Low rainfall has affected the state in this way in the past and has the potential to do so in the future as well. The dependence of hydropower on hydrological conditions means this major source of energy is vulnerable to drought, and may be less reliable if droughts increase in severity or duration into the future.

³³ Carnie, TL, Berry, HL, Blinkhorn, SA and Hart, CR 2011, *In their own words: Young people’s mental health in drought-affected rural and remote NSW*, Australian Journal of Rural Health volume 19.

4. Future Impacts of Drought

4.1. Future Drought Conditions

Under a high emissions scenario it is projected that Tasmania will continue to get hotter in the future³⁴. It is expected that by 2050 Tasmania will see an increase in average annual temperature of between 0.9-1.7 °C.

Hot days (>30 °C) will increase in Hobart from 4-5 days per year to 8 days.

Tasmania can expect longer fire seasons, with around 40% more very high fire danger projected by 2050.

While mean rainfall to 2050 will remain relatively similar, it is expected to occur at different times of the year than it has historically, and in more frequent and severe extreme events.

Drought frequency for the Southern region will remain similar to the existing frequency over the last 100 years. Although given that there will be increase in extreme heat events, there is a potential for droughts to be more intense.

4.2. Future Climate Trends for the Southern Region

Using data from My Climate View³⁵ for 2030, 2050 and 2070, future rainfall, temperature and evapotranspiration trends for four towns in the Southern region were assessed to provide a snapshot of future climate trends across the region.

4.2.1. Rainfall

Mean annual rainfall in the Southern region is projected to stay relatively stable over the next 50 years. However, Figure 4-1 indicates that rainfall on the east coast, central highlands and through the Southern Midlands will start to decrease. The spread across the seasons is also projected to change, see Figure 4-2. Summer rainfall changes for the four identified sites varies from a 5% increase (Bothwell) to an 8% decrease (Huonville), the overall average across the four sites is a decrease of 2% in summer rain. In total, average annual change in rainfall for the four identified sites is a 2% decrease.

Modelling completed by Climate Futures also indicates that there is likely to be an increase in extreme rainfall events. While rainfall may decrease overall, the fewer rainfall events that do occur are likely to be more intense.




Figure 4-1: Expected mean annual rainfall to 2050 (source: [LISTmap](#)).

³⁴ CSIRO, <https://www.climatechangeinaustralia.gov.au/en/changing-climate/state-climate-statements/tasmania/>, accessed 9/8/23

³⁵ Australian Government <https://myclimateview.com.au/>, accessed 9/08/23



Figure 4-2: Mean seasonal rainfall projected changes from current averages to 2070 (source: [My Climate View](#)).

4.2.2. Temperature & Hot days

The average maximum temperature for the key Southern region towns is projected to increase by 21-2.2°C by 2070, with the average minimum temperatures also projected to rise by 2-2.2°C. Hence, there will be less frosts and more hot days. Figure 4-3 shows the projected increase in days over 30°C.




Figure 4-3: Changes in the number of days over 30°C to 2070 (source: *My Climate View*).

4.2.3. Evapotranspiration

Evapotranspiration is projected to increase across the Southern region as shown in Figure 4-3. For summer, the projected average increase in evapotranspiration across the four sites from current to 2070 is 5%. When comparing this to a mean summer rainfall change of -2% of summer rainfall, the indication is that soil moisture levels will generally trend down in summer, which has the potential to affect plant growth, hence irrigation requirements will increase.




Figure 4-4: Mean seasonal evapotranspiration projected changes from current averages to 2070 (source: *My Climate View*).

Water Availability

Catchment modelling by Climate Futures Tasmania in 2010 utilised the CSIRO’s Sustainable Yields models for Tasmanian catchments as well as the IPCCs A2 Scenario climate models to project future catchment water availability for 78 rivers in Tasmania. Both the Department of Natural Resources and Environment’s (NRE) current catchment data, as well as Tasmania Irrigation water availability modelling are based on the CSIROs Sustainable Yields models. Work is currently underway to update future modelling for water availability using CMIP5 and CMIP6 climate data which is produced by the World Climate Research Programme and widely used as the basis for climate projections, however this was not available at the time of writing this report.

For catchments in the Southern region of Tasmania, the Climate Futures for Tasmania Technical Report projected that there will be a variable change in annual flows³⁶. Projected reduced rainfalls in the Central Highlands will reduce flows in catchments that receive runoff from this area.

This will also see a projected reduction of inflows to Lake Crescent/Sorell by up to 23%. However, Lake Tooms is expected to see an increase of inflows by up to 25% and Craigbourne Dam inflows are projected to increase by 24%. Changes to catchments flows have implications for agriculture, especially for availability of irrigation water (both positive and negative).

4.3. Impacts on Agriculture

As the climate warms, the conditions of locations in the Southern region will change. The CSIRO's climate analogues³⁷ provide a snapshot of how towns' future projected climatic conditions will relate to the existing conditions of other towns in Australia to allow climatic comparisons. Figure 4-5 provides a snapshot of three locations in the Southern region (Bridgewater, Cygnet and New Norfolk) and shows the mainland towns that their conditions will be most similar to in 2050. These changes in local Tasmanian conditions will, in all likelihood, impact the types of crops and livestock best suited to be grown in the region. For example, the future climate of New Norfolk will be similar to a current day Cooma, NSW.




Figure 4-5: Climate analogues for three Southern region locations for 2050.

³⁶ Bennett, JC, Ling, FLN, Graham, B, Grose, MR, Corney, SP, White, CJ, Holz, GK, Post, DA, Gaynor, SM and Bindoff, NL 2010, *Climate Futures for Tasmania Technical Report: Water and Catchments*, Antarctic Climate and Ecosystems Cooperative Research Centre, Tasmania.

³⁷ Climate analogues available at: <https://www.climatechangeinaustralia.gov.au/en/projections-tools/climate-analogues/>

With changing climate conditions, agricultural enterprises will become increasingly reliant on irrigation water and natural resource management techniques that assist with retaining as much moisture in the soil as possible. More irrigation schemes will be coming online in the Southern region in the next few years. This will alleviate more producers' sole reliance on adequate rainfall. If agricultural holdings are able to secure reliable access to water, then overall, the productivity of some agricultural crops will remain similar.

Figure 4-6 illustrates modelling of pasture suitability in the Southern region for current, 2030 and 2050 scenarios. In general, it shows that the region will continue to support a wide range pasture species, however it does identify areas within the Central Highlands and Southern Midlands where the species diversity will reduce. Figure 4-7 indicates that the area suitable for cherry production will become more reliant on temperature management by 2050. Figure 4-8 indicates that areas of the Southern region, particularly in the east will become more suited for wine grape production. It is important to note, that the enterprise suitability mapping shown in these figures assumes that the mapped crops will have an adequate supply of irrigation water.





Figure 4-6: Pasture Index Suitability Model³⁸ for 2018, 2030 and 2050 (source: [LISTmap](#)).

³⁸ The Versatility Index provides a scoring system out of 100 that considers the suitability of 10 pasture species. A high score indicates that there are a range of pasture species that can be grown, hence it has a high versatility.



Southern Region: Enterprise Suitability Mapping - Cherries (2018 to 2050)





Figure 4-7: Cherries Suitability Model for 2018, 2030, 2050 (source: [LISTmap](#)).



Southern Region: Enterprise Suitability Mapping - Table Wine Grapes (2018 to 2050)




Figure 4-8: Table Wine Grapes Suitability Model for present, 2023, 2050 (source: [LISTmap](#)).

Case studies: Soil moisture retention in practice

Improving soil health through increased organic matter will improve the amount of moisture able to be retained in agricultural soils and improve the drought resilience of production systems. Below are two Tasmanian examples of soil moisture retention in practice.

Strip-tillage in vegetable production

Vegetable growers can reduce the greenhouse impact of vegetable production by maintaining and preventing further loss of stored soil carbon (mitigation) which will also have soil health and productivity benefits.

Increasing organic matter inputs (crop residues, cover crops and composts) and reducing losses (cultivation, fallow and high disturbance harvesting) are key to maintaining soil carbon stores and improving soil health and productivity.

Strip-till is a system of cultivation that works strips of soil where the crop will be planted or sown and leaves most of the soil covered and undisturbed. Strip tillage is currently used in Tasmania for brassica seedlings, fodder beet and carrot seed.

“We are getting better moisture retention. A crop like brassica prefers the trash on the surface of the soil due to nitrogen hold-up. With a short amount of time to get [ground] ready, logistically strip-till is way better” – Joe Cook, Farmer & Contractor

The main benefits being seen include improved water infiltration and retention, healthier crops, fuel savings, fertiliser placement and efficiency.³⁹

Under vine mulches in viticulture

Soil moisture retention in perennial horticulture production systems, such as viticulture, is critical due to the impact of water on crop quality, high-value crop and large cost of re-planting.

Firstly, it is important to ensure that vineyard irrigation delivery is designed to run efficiently. Effective irrigation design is crucial to ensure that vineyard irrigation systems deliver the quantities of water required in the most efficient manner.

Secondly, irrigation needs can be reduced through using mulches to help retain soil moisture. The use of compost or other organic amendments / mulches under vines will reduce water evaporation and can increase soil organic carbon thus leading to increased water holding capacity, increased nutrient availability and increased microbial activity.

“We mulch prunings into the ground and blow grass mulch under the vines.” – Clover Hill Wines, Tea Tree (Coal River Valley) and Lebrina (Pipers Brook), Tasmania

Other practices that will ‘store’ carbon on farm can be used to improve soil health, such as native revegetation and planting slow-growing inter-row perennial pastures.⁴⁰

³⁹ Soil Wealth Integrated Crop Protection (2019) Strip-till in Tasmania; A reduced till farming system, <https://www.soilwealth.com.au/resources/videos-and-apps/strip-till-in-tasmania-a-reduced-till-farming-system/> and <https://www.soilwealth.com.au/imagesDB/news/StripTillInTasmanianVegetableCrops2018.pdf>

⁴⁰ Tas Farming Futures (2015) Improving efficiency and carbon storage to reduce vineyard costs and GHG emissions; Case study, <https://nre.tas.gov.au/Documents/Clover%20Hill%20Case%20study%20FINAL%20Aug15.pdf>

4.4. Impacts on Natural Environment

The 2030 Natural Resource Management (NRM) Strategy Southern Tasmania Version 3.0⁴¹ lists climate change impacts as one of the four main threats impacting Tasmania's natural environment in the near future (to 2030). The other three threats are habitat loss and fragmentation, biosecurity, and urban and industrial encroachment. Changes in the environment (including drought frequency and severity) will impact the natural environment and ecosystem services that the community relies on. Industry priorities and increasing human population may also place pressures on the natural environment. Possible climate change impacts on the region are detailed in Figure 4-9.

Drought can have significant effects on biosecurity (measures and practices taken to prevent the introduction, establishment, and spread of pests, diseases, and invasive species that could harm the environment, economy, and human health) in Australia due to its impact on ecosystems, agriculture, and the movement of people and goods. Possible effects are:

- **Increased Vulnerability to Invasive Species:** Drought weakens native plants and ecosystems, making them more susceptible to invasion by non-native species. Invasive species can outcompete native species, disrupt ecosystems, and spread diseases that harm both wildlife and agriculture.
- **Altered Pest and Disease Dynamics:** Drought can affect the behaviour and distribution of pests and diseases. Some pests may thrive under drought conditions, while others may become more dormant. This can complicate pest management strategies and lead to unpredictable outbreaks.
- **Changes in Agricultural Practices:** During droughts, agricultural practices may change in response to water scarcity. Farmers might switch to more water-efficient crops or alter their irrigation methods. These changes could inadvertently introduce new pests or diseases to an area.
- **Weakened Quarantine Measures:** Drought can strain resources, including those allocated to biosecurity measures such as inspections at ports and borders. Reduced resources might lead to weakened quarantine efforts, potentially allowing the introduction of new pests and diseases into the country.
- **Livestock and Animal Health:** Drought can impact the health and welfare of livestock and wildlife by reducing the availability of water and nutritious forage. Animals weakened by drought are more susceptible to diseases, and the movement of animals in search of water can facilitate the spread of diseases between regions.
- **Human Behaviour Changes:** Drought may cause people to move in search of better conditions, leading to the unintended movement of pests, diseases, and invasive species. Additionally, economic pressures from drought might lead to illegal or unregulated trade, which can contribute to the spread of biosecurity threats.
- **Altered Ecosystems:** Drought can disrupt ecosystems and biodiversity, leading to changes in plant and animal populations. This can have cascading effects on the interactions between species, potentially favouring the spread of certain pests or diseases.
- **Water Scarcity and Disease Spread:** Water scarcity during drought can lead to water storages becoming stagnant, providing breeding grounds for disease-carrying mosquitoes. This can increase the risk of vector-borne diseases such as dengue fever and Ross River virus, which may have the potential to develop into greater risks in the future. Decreased river flows can lead to a range of detrimental environmental impacts such as increases to nutrient and pollutant concentrations, increases to water temperature, and declines in the extent and condition of riparian vegetation. Water scarcity also impacts the amount and quality of water available for irrigation and agricultural needs.

⁴¹ <https://nrmsouth.org.au/wp-content/uploads/2023/04/NRM-South-Regional-Strategy-2030.pdf>

	Direction	Confidence	Tasmanian Detail
 Temperature	INCREASE ↑	Very High	By 2050, Tasmania is projected to experience an increase of at least 1.5 °C, in all seasons, even if a low-emissions scenario is followed.
 Extreme temperature	INCREASE ↑	Very High	More hot days and warm spells are projected, with fewer frosts. Projections under a high emission scenario indicate an increase from 1.6 days over 35°C to up to 4.2 days, and a decrease in days under 2°C from 9.1 to 0.3 days by 2090. All scenario assessments indicate an increase in all types of high temperature extremes, including maximum daily temperatures, heatwave intensity and heatwave duration.
 Rainfall	DECREASE ↓	Medium	Strong regional differences. A decrease in spring rainfall (10-20% less rain in some areas, and up to 25% by 2050 in the central north). Large reduction in autumn rainfall (up to 50% less rain for some areas relative to 2010-2020). Some regional increases in winter rainfall over the next 20-50 years (e.g. western Tasmania).
 Extreme rainfall	INCREASE ↑	Medium	Increased intensity of extreme rainfall events is projected. Increase in extreme rainfall, particularly along east coast in summer and autumn. Frequency of westerly rain-bearing fronts is expected to decrease gradually, and east-coast lows to increase. Intermittent, more damaging, high intensity rainfall is possible.
Evaporation rate	INCREASE ↑	High	Higher decline in soil moisture during summer and autumn in Tasmania. Increased evaporation associated with warming.
 Drought	INCREASE ↑	Medium	Episodic and regional nature of drought events will continue. Projected decrease in rainfall and increase in evaporation contribute to more time in drought. The east coast of Tasmania will stay especially drought prone.
 Wind speed	INCREASE ↑	Medium	Stronger wind speed in winter in western Tasmania, and a decrease in summer wind speed. Possible increase in extremes.
 Fire weather	INCREASE ↑	High	A harsher fire-weather climate in the future with consistent increases in fire weather projected for Tasmania. A steady increase in fire danger throughout the current century, including an increase in the length of the fire season and an increase in the number of days at the highest range of fire danger. By the end of this century: twice the fire danger, over twice the area, twice as often in Tasmania. This is an eight-fold increase in fire risk.
 Sea level	INCREASE ↑	Very High	Mean sea level will continue to rise and height of extreme sea-level events will also increase. By 2030, between 0.07-0.19 m rise from 1986-2005 sea levels is projected. By 2090, 0.27-0.66 m under low emission scenario and 0.39-0.89 m under high emission scenario. Some exposed locations are projected to see a 1-in-100-year coastal inundation event move towards an event occurring almost every year (during the annual high tide).
 Ocean temperature	INCREASE ↑	High	South-eastern Australia is a hotspot for ocean temperature changes, with projected rise of >3°C under a high emission scenario. The western Tasman Sea is considered a global ocean warming hotspot. By 2060, intense marine heatwave events are expected to increase.
 Ocean acidification	INCREASE ↑	Medium	Benthic and pelagic calcifiers, such as diatoms, molluscs and deep water coral, will show reduced calcification rates and/ or increased dissolution.

Figure 4-9: Summary of climate change impacts in Tasmania (reproduced under Creative Commons licence 3.0 from 2030 NRM Strategy Southern Tasmania⁴²).

⁴² <https://nrmsouth.org.au/wp-content/uploads/2023/04/NRM-South-Regional-Strategy-2030.pdf>

4.5. Impacts on Communities

While future droughts in the Southern region of Tasmania are not currently projected to be more frequent they may be more severe than in the past. This may have significant implications for the community, particularly given that other extreme events such as rainfall and heat are projected to increase. These impacts are likely to lead to more severe flood and bushfire events.

The recent history on the eastern seaboard of mainland Australia is testament to the likely increase in extreme events. 2017-2019 saw many areas in drought, followed by the Black Summer fires of 2019/20, the pandemic of 2020, and two years of severe flooding.

Increases in extreme weather events have significant implications for agriculture and the communities where agriculture occurs. The resilience of these communities will be tested.

Agriculture, forestry and fishing is currently not a high employer in the region (3.5%). However, extreme events that impact on the agricultural industry have the potential to affect a large number of residents in southern Tasmania who rely on the agricultural industry for employment⁴³. This is particularly significant for Southern Midlands where 29% of employed people are employed in the agriculture, forestry and fisheries sector, as well as Huon (24%), Glamorgan-Spring Bay (18%) and Tasman (18%).

⁴³ National Institute of Economic and Industry Research 2023, *Northern Tasmania: Employment by industry (total)*, Economy: Informed Decisions, accessed 10/10/2023 from <<https://economy.id.com.au/tasmania/employment-by-industry?WebID=420&sEndYear=2021>>

5. Risk, Adaptive Capacity & Resilience

5.1. Drought Impact (Risk)

ABARES Community Vulnerability & Resilience to Drought Index (CVRDI) has assessed the potential drought impact (risk) for all LGAs across Australia. This was completed by using the method shown in Figure 1-1. The index provides a score of between 0 and 1, with 0 meaning lower risk and 1 meaning higher risk. The scores for the LGAs in the Southern region are displayed in Figure 5-1. The LGA with the highest risk ranking is Central Highlands with 0.33. Based on the CVRDI the potential vulnerability of LGAs in the Southern region to drought impacts are on the lower end of the index.




Figure 5-1: Potential drought impact on Southern region LGAs.

5.2. Adaptive Capacity

5.2.1. Framework and Indicators

The adaptive capacity of the Southern region has been assessed utilising data to define the five capitals (physical, natural, financial, human and social). The methodology for this assessment is further discussed in Section 1.5 and Appendix 1. The five capitals have been assessed at the LGA level and are⁴⁴:

- **Physical capital** – items produced by economic activity from other types of capital that can include infrastructure, equipment and improvements in genetic resources (crops, livestock)
- **Natural capital** – the productivity of land, and actions to sustain productivity, as well the water and biological resources from which rural livelihoods are derived
- **Financial capital** – the level, variability and diversity of income sources, and access to other financial resources (credit and savings) that together contribute to wealth
- **Human capital** – the skills, health and education of individuals that contribute to the productivity of labour and capacity to manage land
- **Social capital** – reciprocal claims on others by virtue of social relationships, the close bonds that facilitate cooperative action and the social bridging, and linking via which ideas and resources are accessed.

Two relevant indicators for each capital have been identified that provide data at the LGA level for each capital, recognising that measuring human and social capital is inherently complex.

Table 5-1: Indicators used for each capital.

Capital	Indicator 1	Indicator 2
Physical	Megalitres (ML) Irrigation Scheme water currently available per ha of agricultural land	ML of stored water (irrigation & stock) per ha of agricultural land – using NRE registered dam data
Natural	Area of Agricultural land – ABS Census data	Diversity of agricultural enterprises – ABS Census data
Financial	SEIFA Index of Relative Socio-economic Disadvantage	SEIFA Index of Economic Resources
Human	SEIFA Index of Education & Occupation	People with a mental health condition
Social	Percentage of people who volunteer	Percentage of people who have moved (over census period)

The results for each LGA are displayed in Table 5-2 below, and discussed in Section 5.2.5 to Section 5.2.4. All references to census data are to 2021 ABS data, unless otherwise stated.

5.2.2. Physical Capital

The adaptive capacity for physical capital was assessed by the volume of water available via an active irrigation scheme and the volume of water stored in farm storages (for irrigation, stock and aquaculture) over the LGA's total agricultural area.

Within the Southern region there are six gazetted irrigation schemes that provide approximately 31,000ML of irrigation water (Figure 5-2). There is also the Southern Midlands irrigation scheme under development which will provide up to an additional 18,000ML of irrigation water in the coming years. Only existing irrigation schemes have been included in the assessment. However, it is important to note that as the new

⁴⁴ Nelson et al (2007) The potential to map the adaptive capacity of Australian land managers for NRM policy using ABS data, Natural Heritage Trust

schemes come online the physical capital of the associated LGAs will improve.

It is also recognised that there is a cost associated with sourcing water from an irrigation scheme, which has the potential to impact on a land manager’s financial status. The development of irrigation schemes can also impact on the environment. Cost benefits analyses are conducted to determine a scheme’s financial viability, potential community benefits, as well as demonstrating that the scheme is environmentally sustainable prior to development, which is based on sustainable yields⁴⁵.

Volume of farm storage was taken from dam data provided by NRE in August 2023. All registered irrigation, aquaculture and stock dams have been included to understand the overall storage capacity of each region, this has then been divided by ha of agricultural land to give a ML/ha figure for each LGA. Clarence has the highest ML/ha ratio of stored water at 0.46ML/ha of regional councils, while Derwent Valley has the least at 0.02ML/ha. It is noted that many irrigators in the Derwent Valley have summer take allocations from the Derwent River, these were not included in the assessment.




Figure 5-2: Active irrigation schemes in the Southern region.

⁴⁵ <https://www.tasmanianirrigation.com.au/about-tasmanian-irrigation>

5.2.3. Natural Capital

There are many ways to measure natural capital including agricultural land use (as used in this report) as well as biophysical indicators such as ecosystem health and diversity, and native vegetation coverage. From a land management perspective, utilising the land for a diversity of agricultural enterprises can assist with providing greater resilience to climatic factors especially if the mixed uses have differing climatic tolerances (e.g. water use). In general, the Southern region has a high diversity of agricultural enterprises. Glamorgan-Spring Bay Sorell and Clarence have the greatest diversity of agricultural enterprises (39, 33 and 33) respectively). Unsurprisingly, Hobart and Glenorchy have the lowest diversity of agricultural enterprises, with 1 and 2 respectively. These two LGAs also have the least area of agricultural land.

In addition, sustainable agricultural practices that improve soil health, protect waterways and increase vegetation coverage are important to assist with increasing natural capital.

5.2.4. Financial Capital

The Socio-Economic Indexes for Areas (SEIFA) Index of Relative Social Disadvantage (IRSD) is widely quoted as an indicator of community socio-economic disadvantage. The Index of Economic Resources (IER) has a greater focus on financial variables and includes 'variables that correlate to high or low wealth as well as variables that are indicators of high or low income'.⁴⁶

The Southern region LGAs in descending order of disadvantage (most to least), and with Tasmanian IRSD decile ranking in brackets, are: Brighton and Central Highlands (both 2), Derwent Valley (3), Glenorchy and Tasman (both 4), Glamorgan-Spring Bay and Southern Midlands (both 6), Huon Valley (8), Sorell (9) and Hobart and Clarence (both 10)

The IER decile rankings are very similar but with Brighton and Glenorchy showing the lowest economic resources.

5.2.5. Human Capital

Diagnosed mental health condition/disorder rates across the region range from 9.65% (Glamorgan-Spring Bay) to 13.34% (Derwent Valley). The Tasmania average is 11.45%, while the national average is 8.78%. The figures indicate much of the Southern region and Tasmania as a whole is well above the national average for diagnosed mental health conditions.

Brighton demonstrates a consistently low score across all SEIFA indexes, sitting in the 2nd decile for Tasmania. Together these scores reflect an area of disadvantage. Brighton also is in the 2nd Tasmanian decile for the IEO, Central Highlands the 3rd. While, Clarence, Hobart and Kingborough have a high level of education and occupation (in the 10th decile).

5.2.6. Social Capital

The measures identified for social capital were the percentage of residents in an LGA who volunteered and the percentage of people who have moved address in the last five years. For the Southern region, interestingly Hobart has the highest level of volunteerism at 25.2% and also has the highest level of people movement (45.36%). This bucks the trend from the north and north west of Tasmania, where the main population centres generally had the lowest level of volunteerism and the highest level of people movement. The regional community of Tasman has a moderately high level of volunteerism at 22.2% and a lower people movement level at 32.67%. This indicates that this community is a socially connected community, that sees a greater percentage of people involved in the community and with less movement within and out of the community. Brighton has the lowest level of volunteerism at 10.4%. The Tasmanian average across LGAs for volunteerism is 18.8% and people movement is 35.34%.

⁴⁶ <https://www.treasury.tas.gov.au/Documents/Discussion%20Paper%20DP19-01%20-%20SOCIO%20ECONOMIC%20FACTORS%20IN%20THE%20BASE%20GRANT%20MODEL%20-%20Proposal%20to%20replace%20the%20Unemployment%20Cost%20Adjustor%20with%20a%20SEIFA%20based%20Cost%20Adjustor.pdf>

Table 5-2: Five capital indicators data for Southern region LGAs.

LGA	Physical		Natural		Financial		Human		Social	
	ML of storage per ha. ag land	ML irrigation scheme water per ha. ag land	Area of Ag Land (ha) [^]	Diversity of Ag Enterprises (number)	SIEFA-IRSD*	SIEFA-IER*	SIEFA-IEO*	Mental Health	People Movement	Volunteerism %
Brighton	0.03	0.39	4,596.20	22	2	2	2	13.20%	35.40%	10.4%
Central Highlands	0.06	0.14	164,188.72	26	2	5	3	10.10%	34.57%	15.3%
Clarence	0.38	0.46	8,899.04	33	10	9	10	11.27%	35.95%	17.5%
Derwent Valley	-	0.02	19,195.31	21	3	5	2	13.34%	31.59%	12.7%
Glamorgan-Spring Bay	0.02	0.17	93,916.07	39	6	6	7	9.65%	36.90%	21.7%
Glenorchy	-	3.15	23.43	2	4	2	6	12.65%	39.32%	14.2%
Hobart	-	257.37	1.14	1	10	7	10	10.53%	45.36%	25.2%
Huon Valley	-	0.23	15,456.69	20	8	8	9	12.22%	35.45%	19.0%
Kingborough	-	0.10	9,229.06	17	10	10	10	10.97%	36.97%	23.5%
Sorell	0.14	0.42	13,691.20	33	9	9	8	12.10%	37.82%	15.5%
Southern Midlands	0.05	0.29	179,594.56	28	6	10	4	10.92%	28.00%	15.4%
Tasman	-	0.21	2,570.40	11	4	4	7	12.94%	32.67%	22.2%

*See Appendix 2 for explanation of SEIFA indexes and ranking system.

[^] rounded to 2 significant figures

5.2.7. Results

Figure 5-3 shows a visual representation of the assessed indicative adaptive capacity to drought of the LGAs in the Southern region once they have been ranked relative to other Tasmanian LGAs (as described in Appendix 1). As shown below Clarence displays higher adaptive capacity to drought, whereas Central Highlands and Derwent Valley display lower adaptive capacity to drought.

It is important to note that adaptive capacity will be reduced by more frequent, longer duration and severe drought events.




Figure 5-3: Adaptive capacity to drought of Southern region LGAs.

5.3. Resilience to Drought

To understand the relative resilience of communities in the Southern region to drought, the results from the adaptive capacity assessment have been combined with the CVRDI to assess the resilience score for each LGA. A rating of 0 means the LGA is considered to have a higher resilience and 1 means the LGA is considered to have a lower resilience to drought. The scores displayed in Figure 5-4 suggest that Central Highlands, Glamorgan-Spring Bay and Tasman LGAs have lower resilience to drought and Kingborough, Sorell and Clarence LGAs have higher resilience to drought and are therefore less vulnerable.




Figure 5-4: Drought resilience of Southern region LGAs.

5.4. Gaps in Preparedness

The identified long-term outcomes for the Drought Resilience Plans include:

- More primary producers preserve natural capital while also improving productivity and profitability
- Stronger connectedness and greater social capital within communities, contributing to wellbeing and security.
- Communities implement transformative activities that improve their resilience to drought

When identifying gaps in preparedness they need to be linked back to the gaps that are going to assist with achieving the above outcomes. Resilience is the capacity of a system to cope with change and continue to evolve in positive ways. Figure 5-5 shows that efforts that influence a system are most effective when intervention occurs at a transformative, deeper level. When systems persist and operate 'above the water line' of the iceberg, and respond to the shocks and events they can see or easily foresee, they will be less resilient. When systems are reactive, they fail to address the underlying patterns, processes and systemic structures that lead to those shocks and events. Hence, to address any gaps in the preparedness to future drought (and other extreme events) in the long-term the Regional Drought Resilience Plan should develop actions and solutions that will assist communities and land managers to adapt and transform over a period of time.




Figure 5-5: Iceberg model of systemic change⁴⁷.

Based on the data and on anecdotal evidence some of the gaps in preparedness that have been identified through this report are outlined in the table below.

⁴⁷ RMCG & Australian Resilience Centre (2020) *Goulburn Murray Resilience Strategy*

Table 5-3: Gaps in drought preparedness in the Southern region.

Theme	Gap
Agriculture and industry	<ul style="list-style-type: none"> • Technology literacy. There is a range of technology available to assist land managers to make short- and long-term decisions based on seasonal forecast information. However, there is often a gap between the available technology and land managers either knowing it exists or being able to utilise it. • Awareness of climate change. General awareness of how the Tasmanian climate is changing, what this may mean for local communities and how it will impact agricultural enterprises into the future. • Diversity of agricultural enterprises. Lack of diversity of agricultural enterprises in some areas will mean these areas are less resilient to cope with future drought scenarios. • Biosecurity. To address drought related biosecurity challenges, Australian biosecurity authorities need to adapt their biosecurity strategies for drought periods. This may involve increased monitoring for invasive species, maintaining robust quarantine measures, providing guidance to farmers on best practices for pest and disease management during drought, and educating the public about the risks of introducing and spreading biosecurity threats during difficult environmental conditions.
Natural environment	<ul style="list-style-type: none"> • Adoption of sustainable land management. There is a lack of data available around the uptake of NRM practices on farms at an LGA or lower level. Being able to better track this data would assist in understanding how individual landowners and LGAs have developed natural responses to assist with mitigating drought impacts.
Communities	<ul style="list-style-type: none"> • Diagnosed mental health disorders. Diagnosed mental health disorder rates across Tasmania in general are much higher than the national average. It is important to note that this data does not include undiagnosed conditions, hence the rates are likely to be much higher. Mental health is an important factor when understanding individual resilience to crisis. • Undiagnosed mental health disorders. It is noted that the mental health indicator used for human capital only includes people with a diagnosed condition. Hence, the actual percentage of people with undiagnosed conditions is likely to be much higher, particularly in regional communities. This will need to be factored into any future actions that may look to provide community support for mental health.
Data access and limitations	<ul style="list-style-type: none"> • Climate data. Information on long term climate modelling for Tasmania is spread across a number of different agencies, source and platforms, which can make it difficult for individuals to find the most up to date and relevant information for the region and state in general. Better consolidation of data sources at a government level would assist land managers greatly. • Irrigation cease to take data. There is little irrigation cease to take data available for the regions, and none is publicly available prior to the 2018/19 irrigation season. Making more of this data available would assist in being able to better understand how dry spells have influenced land managers ability to irrigate in the past, and how they may be impacted by future dry periods. • Past drought impact data. Data is lacking on how specific previous droughts have impacted individuals and communities in Tasmania. While it is known that extreme weather events have health impacts (mental and physical) and financial impacts, providing specific examples was difficult. It is noted that this information is often considered sensitive and this likely would be a key reason for why this data is not publicly available. However, this type of data would also be beneficial in that it could further assist with identifying the most vulnerable communities within the region⁴⁸.

⁴⁸ Note: the University of Tasmania Menzies Wicking Centre could potentially supply data and assist with filling some of the identified gaps.

6. Building Resilience to Drought

6.1. Key Observations and Themes

Droughts occurred in the Southern region of Tasmania in 2006, 2008, 2015 and 2019. These events offer insights on the impacts to agriculture and the natural environment, and the potential resilience of communities to future droughts.

Drought resilience was determined by analysing the potential drought impact (risk) and adaptive capacity of each of the twelve LGAs in the Southern region. This showed Clarence had higher adaptive capacity potential to drought, whereas Central Highlands and Derwent Valley display lower adaptive capacity to drought. The potential drought impact (risk) showed Central Highlands with higher risk rating, while Kingborough was lower based on the ABARES Community Vulnerability & Resilience to Drought Index (CVRDI). The potential vulnerability of LGAs in the Southern region to drought impacts are on the lower end of the index.

Therefore, the Southern region has moderate resilience to manage future drought conditions. Central Highlands, Glamorgan-Spring Bay and Tasman LGAs have lower resilience to drought and Kingborough, Sorell and Clarence have higher resilience to drought. However, it is important to note that climate change will increase the frequency, severity and duration of extreme events such as periods of intense heat and rainfall. While the drought risk may be projected to moderately increase in most of the Southern region, it is going to be one of many factors that land managers and communities need to prepare for and respond to in the future.

While agriculture, forestry, and fisheries are not major drivers of the economy in the Southern region in terms of total employment, they are important drivers for individual councils (such as Huon Valley and Southern Midlands), where they employ a higher number of people. As the climate continues to change these industries need to continue to adapt and transform to ensure they are resilient to drought and other changes in climate.

Climate change is already impacting on agriculture and the natural environment and communities on which it relies. If communities can increase their adaptive capacity and resilience to future drought events, then it will also assist in increasing their resilience to other extreme events. It is important to note that more frequent, longer duration and severe droughts may reduce adaptive capacity.

Much work has been done to provide secure water supplies for agriculture in the region, through irrigation schemes and individual landowner investments in storage dams. In general, the region also has a good diversity of agricultural enterprises, both at the property and regional levels. These are two important factors that assist with the region's existing resilience to drought. The development of the Regional Drought Resilience Plans will help to identify regional needs, priorities and challenges and inform future investment to improve economic, social and environmental resilience to drought.

6.2. Recommendations

Based on the key findings in this report the following recommendations have been identified.

Table 6-1: Recommendations.

Theme	Recommendation
Use this report to inform community engagement undertaken by the Regional Project Coordinator	<ol style="list-style-type: none"> 1. Test the risk, adaptive capacity and resilience to drought findings by LGA with the community to see if it reflects on-ground experience to past events. 2. Investigate the My Climate View forecast data as a tool to support further community engagement, which explores future climate predictions for individual towns and provides a specific snapshot of how conditions will change in the coming years.
Utilise the data in this report to inform the development of the Regional Drought Resilience Plan	<ol style="list-style-type: none"> 3. Undertake win-win, no regrets actions to assist with short to medium-term adaptation to future drought conditions (i.e. avoid maladaptation). 4. Ensure the Regional Drought Resilience Plan develops long-term transformative actions that consider all five capitals; physical, natural, financial, human and social. 5. Prioritise action in those communities with lower resilience by building adaptive capacity and reducing vulnerability to potential impacts of drought. This includes Central Highlands, Glamorgan-Spring Bay and Tasman LGAs in the Southern region. 6. Continue to develop irrigation schemes where feasible that balance environmental water needs. This is a priority in the Southern Midlands area. 7. Work with land managers to continue to improve and diversify their agricultural operations to be more adaptable to changing climatic conditions as well as extreme climate events (such as drought). This will have broader benefits for agriculture, the natural environment and communities. For example, this may include sustainable agriculture practices that improve soil health through increasing organic matter inputs and reducing losses for greater soil moisture retention. 8. Build technical literacy in regions to enable land managers to utilise current and emerging technology to better plan and prepare for changes in seasonal conditions. 9. Ensure actions consider community health, including mental health, as important aspects of resilience in the Southern region. Be aware that rates of mental health disorders are likely to be higher than are reported.
Undertake monitoring, evaluation, reporting and learning (MERL) for the Regional Drought Resilience Plan	<ol style="list-style-type: none"> 10. Establish clear, measurable and robust indicators of drought resilience in the Regional Drought Resilience Plan, informed by this report and emerging best-practice research. 11. Monitor drought resilience over time and update the Regional Drought Resilience Plan as required, including supporting data.

References

Alston, M and Kent, J 2004, *Social impacts of drought: a report to NSW Agriculture*, Centre for Rural Social Research, Charles Sturt University.

Australian Bureau of Agricultural and Resources Economics and Sciences (ABARES) *Community Vulnerability & Resilience to Drought Index*. https://www.agriculture.gov.au/abares/research-topics/climate/drought/resilience#community-vulnerability-and-resilience-to-drought-index-cvr-di_2 accessed August 2023

Australian Bureau of Statistics (ABS). *Estimated Value of Agricultural Operations through the Agricultural Commodity Survey*. <https://www.abs.gov.au/statistics/industry/agriculture/value-agricultural-commodities-produced-australia/2021-22> accessed August 2023

Australian Bureau of Statistics (ABS). *Socio-Economic Indexes for Areas (SEIFA)*. <https://www.abs.gov.au/websitedbs/censushome.nsf/home/seifa> accessed August 2023

Australian Standard, 2015. *ISO 14001:2015 Environmental Management*. Australian Standard

ACE CRE (2010). *Climate Futures for Tasmania – Water and Catchments Summary*. The Antarctic Climate & Ecosystems Cooperative Research Centre, Hobart, Tasmania

ACE CRE (2010). *Climate Futures for Tasmania – Impacts on Agriculture Summary*. The Antarctic Climate & Ecosystems Cooperative Research Centre, Hobart, Tasmania

ACE CRE (2010). *Climate Futures for Tasmania – General Climate Impacts Summary*. The Antarctic Climate & Ecosystems Cooperative Research Centre, Hobart, Tasmania

Bennett, JC, Ling, FLN, Graham, B, Grose, MR, Corney, SP, White, CJ, Holz, GK, Post, DA, Gaynor, SM and Bindoff, NL 2010, *Climate Futures for Tasmania Technical Report: Water and Catchments*, Antarctic Climate and Ecosystems Cooperative Research Centre, Tasmania.

Bureau of Meteorology (BoM). *Understanding Drought*. <http://www.bom.gov.au/climate/drought/knowledge-centre/understanding.shtml>, accessed August 2023

Bureau of Meteorology BoM). *Historical Rainfall Maps for Tasmania*. <http://www.bom.gov.au/climate/maps/rainfall/> accessed August 2023

Caldwell, K and Boyd, CP 2009, *Coping and resilience in farming families affected by drought*, Rural and Remote Health volume 9.

Carnie, TL, Berry, HL, Blinkhorn, SA and Hart, CR 2011, *In their own words: Young people's mental health in drought-affected rural and remote NSW*, Australian Journal of Rural Health volume 19.

Climate Change Australia. *Tasmania Changing Climate*. <https://www.climatechangeinaustralia.gov.au/en/changing-climate/state-climate-statements/tasmania/> accessed August 2023

Climate Change Australia. *Climate Analogues*. <https://www.climatechangeinaustralia.gov.au/en/projections-tools/climate-analogues/> accessed August 2023

CSIRO. *My Climate View*. <https://myclimateview.com.au/> accessed August 2023

Department of Agriculture, Water and the Environment (2020) Future Drought Fund (Drought Resilience Funding Plan 2020 to 2024) Determination 2020, https://ehq-production-australia.s3.ap-southeast-2.amazonaws.com/a69d99c1b753c9e93cdf88ce1fd6d723997f78d9/original/1584927882/Drought_Resilience_Funding_Plan.pdf_7c8e152b3f40ab97f9e17b39fc4ce42a?X-Amz-Algorithm=AWS4-HMAC-SHA256&X-Amz-Credential=AKIA4KKNQAKIOR7VAOP4%2F20231016%2Fap-southeast-2%2Fs3%2Faws4_request&X-Amz-Date=20231016T023846Z&X-Amz-Expires=300&X-Amz-SignedHeaders=host&X-Amz-Signature=20de2119787c15ed7867f67c210e8f670c91122beb19b9d4522b7c3014993942

Department of Climate Change, Energy, the Environment and Water. *Science, Maps and Data*. <https://www.dcceew.gov.au/environment/land/nrs/science> accessed August 2023

Department of Natural Resources and Environment Tasmania (NRE). LISTmap Climate Data. <https://maps.thelist.tas.gov.au/listmap/app/list/map> accessed August 2023

Department of Natural Resources and Environment Tasmania (NRE). *Annual River Reports*. Available at: <https://nre.tas.gov.au/water/water-data/annual-river-reports>

Department of Natural Resources and Environment Tasmania (NRE). *Tasmania's Water Catchments*. <https://nre.tas.gov.au/water/a-guide-to-water-in-tasmania/tasmanias-water-catchmentstasmanian> accessed August 2023

Department of Primary Industries, Parks, Water and Environment Resource Management and Conservation Division 2010, *Vulnerability of Tasmania's Natural Environment to Climate Change: An Overview*, Tasmanian State Government.

Edwards, B, Gray, M and Hunter, B 2008, *Social impacts associated with drought*, Australian Institute of Family Studies, Australian Government.

Fleming-Muñoz, D.A., Whitten, S. and Bonnett, G.D 2023, *The economics of drought: A review of impacts and costs*, Australian Journal of Agricultural and Resource Economics

Grant, R. (2015) *Extreme Heat Causing Ginger Syndrome in Eucalypts*. ABC Rural, published 26/02/2015. Available at: <https://www.abc.net.au/news/rural/2015-02-26/ginger-tree-syndrome/6263118>

Hooper, S, Ashton, D, Crooks, S, Mackinnon, D, Nicols, P, & Phillips, P 2008, *Farm Financial Performance: Australian Farm Income and Drought Recovery, 2005-06, 2006-07 and 2007-08*, Australian Commodities: Forecasts and Issues.

Hughes, N, Galeano, D and Hatfield-Dodds, S 2019, *Analysis of the effects of drought and climate variability on Australian Farms*, ABARES Insights Issue 6, Department of Agriculture, Australian Government.

Hughes, N, Soh, WY, Boulton, C and Lawson, K 2022, *Defining drought from the perspective of Australian farmers*, Climate Risk Management, Volume 35

International Panel on Climate Change (IPCC), 2012, Glossary or term, In: *Managing Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*. (Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G-K Plattner, S.K. Allen, M. Tignor and P.M. Midley (eds.)). A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC). Cambridge University Press, Cambridge UK, and New York, NY, USA pp. 555-564

International Plant Protection Convention Secretariat 2021, *Scientific Review of the impact of climate change on plant pests – A global challenge to prevent and mitigate plant pest risks in agriculture, forestry and ecosystems*, Food and Agriculture Organisation of the United Nations.

Kancans, R. & Stenekes, N. (2022). Community vulnerability and resilience to drought index – stage 1. ABARES, available at: <https://www.agriculture.gov.au/abares/research-topics/climate/drought/resilience#reports>

Lester, L, Flatau, P and Kyron, M 2022, *Understanding the Social Impacts of Drought*, Centre for Social Impact, University of Western Australia.

Luong, TT, Handley, T, Austin, EK, Kiem, AS, Rich, JL and Kelly, B 2021, *New Insights Into the Relationship Between Drought and Mental Health Emerging from the Australian Rural Mental Health Study*, *Frontiers in Psychiatry* volume 12.

National Institute of Economic and Industry Research 2023, *Northern Tasmania: Employment by industry (total)*, *Economy: Informed Decisions*, accessed 10/10/2023 from <<https://economy.id.com.au/tasmania/employment-by-industry?WebID=400>>

Natural Resource Management South (NRM South). *2030 NRM Strategy for Southern Tasmania*. Available at: <https://nrmsouth.org.au/about-us/our-strategy/#:~:text=Regional%20Strategy%20to%202030%20for%20Southern%20Tasmania&text=Developed%20collaboratively%20across%20all%20three,use%20to%20prioritise%20their%20investment>.

Nelson et al (2007) The potential to map the adaptive capacity of Australian land managers for NRM policy using ABS data, Natural Heritage Trust

Profile ID. *Tasmanian Profile*. <https://profile.id.com.au/tasmania> accessed August 20213

Queensland Government. *The Long Paddock*. <https://www.longpaddock.qld.gov.au/> accessed August 2023

RM Consulting Group (RMCG) & Australian Resilience Centre (2020). Goulburn Murray Resilience Strategy. Available at: <https://www.rdv.vic.gov.au/resources/resilience>

Tasmania Irrigation. *Active Schemes List*. <https://www.tasmanianirrigation.com.au/active-schemes-map> accessed August 2023

Tasmania Irrigation. *Projects Under Development List*. <https://www.tasmanianirrigation.com.au/future-irrigation> accessed August 2023

White, D. H. & Walcott, J.J (2009). The role of seasonal indices in monitoring and assessing agricultural and other droughts: a review. CSIRO Publishing, *Crop & Pasture Science* 60: 599-616

Williams, KJ, Hunter, B, Schmidt, RK, Woodward, E and Cresswell, ID 2021, Land: Soil. In: *Australia State of the Environment 2021*, Department of Agriculture, Water and the Environment, Australian Government.

Appendix 1: Adaptive capacity and vulnerability assessment methodology

1 Introduction

A key requirement of the regional drought data reports is providing data that can assist the regional drought coordinators in determining the vulnerable areas within their regions and to then direct resources where they may be most needed. The general method for determining the vulnerability to drought is shown in the drought vulnerability and resilience conceptual framework (Figure 1). This model is from the ABARES website and is the model that has been adopted for the Future Drought Fund.




Figure A1-1: Drought Risk, Adaptive Capacity and Resilience Model⁴⁹.

This methodology details how we applied this framework in assessing the drought vulnerability and resilience for the Tasmanian regional drought data reports.

2 Determining Adaptive Capacity

Two measures were identified for each type of capital which enabled a consistent and robust measurement of adaptive capacity with available data at the LGA level. All quoted census data refers to 2021, which enables the indicators to be updated with future census data.

2.1 Physical Capital

In looking at physical capital and adaptive capacity the measures identified were the volume of irrigation scheme water per LGA and the volume of farm storages per LGA. To ensure that this was comparable with

⁴⁹ <https://www.agriculture.gov.au/abares/research-topics/climate/drought/resilience#reports>

the amount of agricultural activity per LGA these measures were overlaid by the area of agriculture in each LGA recorded in the latest census data.

The physical capital measures used for each LGA then become:

- **Volume of irrigation scheme water/agricultural area**
- **Volume of farm storage/agricultural area**

2.2 Natural Capital

The measures identified for inclusion in the assessment of natural capital were **areas of agricultural land per LGA** and the **diversity of enterprises**. The area of agricultural land is from the latest ABS census data. The diversity of enterprises was measured by counting the number of commodities with an income in the latest census data. This was at the specific commodity level rather than industry (i.e. broccoli vs vegetables). One gap in the ABS census data is that it only captures agricultural enterprises with a gross annual income of greater than \$40,000. This means that many 'small scale' producers that assist with providing greater diversity of agricultural activities may not be captured in the data.

2.3 Financial Capital

The measures identified for financial capital were drawn from the Socio-Economic Indexes for Areas (SEIFA) indexes – a suite of four indexes. The **Index of Economic Resources (IER)** and the **Index of Relative Socio-Economic Disadvantage (IRSD)** were used from the SIEFA data.

2.4 Human Capital

The measures identified for human capital were the **SEIFA education and occupation index (IEO)** and the **reported mental health data from the latest census**. The only minor caveat with the mental health data set is that it relies on the respondent having been diagnosed for one of the conditions in the census. This may mean that more rural areas with reduced access to mental health services may be under-represented in the data.

2.5 Social Capital

The measures identified for social capital were the **percentage of residents in an LGA who volunteered** and the **percentage of people who have moved address in the last five years**. The people who moved address data includes all people who moved within an LGA or outside of an LGA in the assessed five year period, this data indicates residential mobility (or stability) within an area. People who moved multiple times are only captured once in the data.

It is noted that there are a range of potential social indicators that could be used and that measuring social capital is inherently difficult and imperfect.

2.6 Combining the Measures

These 10 measures were brought together to determine an adaptive capacity score for each LGA. Firstly each LGA was ranked for each individual measure. The two relevant measures were then pooled to give a combined ranking by capital type. Each LGA was then given a percentile rating for each capital type. The five capital types were then brought together to give a total score. Based on this total, each LGA was then given a percentile rating as an adaptive capacity score. As with the ABARES data discussed in section 3, the closer the rating to 1, the more likely the LGA is to experience adverse drought impacts (i.e. the LGA has a lower adaptive capacity to handle drought or other shocks).

3 Determining Drought Impact (Risk)

The ABARES data set Community Vulnerability and Resilience to Drought Index (CVRDI)⁵⁰ has been used as a measure for the drought impact risk (corresponding to 'potential drought impact' in Figure 1). To align with the methodology of scoring the adaptive capacity, the CVRDI data was pulled for Tasmania and then the scores ranked by percentile for Tasmanian Local Government Areas (LGAs). As for the ABARES data set, the closer the rating is to 1, the more at risk the LGA is to drought.

4 Determining Drought Vulnerability & Resilience

As described in section 2.6 and section 3 of this Appendix we then had a Tasmanian percentile rating for each LGA for both drought impact and adaptive capacity. These two ratings were then added together to create a drought vulnerability and resilience score. To ensure this was to the same scale as the drought impact and adaptive capacity ratings, these two scores were averaged together to achieve a final combined drought vulnerability & resilience score. As with sections 2.6 and 3 the closer the rating to 1, the more vulnerable the LGA is to drought.

⁵⁰ <https://www.agriculture.gov.au/abares/research-topics/climate/drought/resilience#reports>

Appendix 2: Socio-economic data (SEIFA)

A key data source that has been used to assist with providing an indicative assessment of the regions' adaptive capacity to drought are the Socio-Economic Indexes for Areas (SEIFA). SEIFA are weighted calculations based on multiple Census variables. These indexes allow comparison between areas, but not necessarily over time, as to possible vulnerabilities and measures of potential resilience such as family structure, education and home ownership. There are four indexes, three of which were used in the Adaptive Capacity assessment:

- The Index of Relative Socio-economic Disadvantage (IRSD) – indicator for Financial Capital
- The Index of Relative Socio-economic Advantage and Disadvantage (IRSAD)
- The Index of Education and Occupation (IEO) – indicator for Human Capital
- The Index of Economic Resources (IER) – indicator Financial Capital

These indexes are described in Table A2-1

The most commonly quoted index is the Index of Relative Socio-economic Disadvantage.

Each of the indexes is built to have 1000 as its midpoint (the mean) and one standard deviation is 100. These are based on Census SA1 areas, the smallest Census data area, from across Australia. LGA scores are population weighted averages of the SA1 areas within that LGA.

Areas within Tasmania have been ranked from lowest to highest score, with the lowest-scoring 10 percent of areas given a decile number of one, up to the highest ten percent which is given a decile number of 10. Each decile represents a population of the same size and this allows meaningful comparisons between areas within the state. Data not shown here, but included in the discussion are percentiles, where similar sized populations are ranked between 1 and 100, for both within Australia and within Tasmania.

SEIFA data for the Southern region LGAs is included in Table A2-2. The individual scores for each index allow comparison with the rest of Australia. All scores are below 1000 (with one exception), indicating that some Southern region LGAs show measures of social disadvantage and relative lack of advantage when compared to other Australian LGAs.

Table A2-1: SEIFA indexes and characteristics⁵¹.

	Index of			
	Relative Socio-economic Disadvantage	Relative Socio-economic Advantage and Disadvantage	Education and Occupation	Economic Resources
Focus	Disadvantage	Advantage and disadvantage	Advantage and disadvantage	Advantage and disadvantage
Summarises Census variables of	Economic and social conditions of people and households – only looks at disadvantage	As for IRSD but looks at advantage and disadvantage	Educational and occupational level (but not income)	Income and housing
No. of variables	15	23	8	14
Types of variables	Low income, low education, high unemployment and unskilled occupations, family structure, disability	Income, internet connection, occupation and education, levels of rent, family structure, disability	Qualifications achieved, current further education, occupations with high/low skill levels, unemployment	Household income, ownership, composition, and levels of rent and mortgage, , unemployment, ownership of an unincorporated enterprise
Most highly correlated variable	Household equivalised income <\$26,000 per year	Disadvantage - Highest education attained is Year 11 or lower Advantage - Household equivalised income >\$91,000 per year	Disadvantage - Highest education attained is Year 11 or lower Advantage – working in a Skill Level 1 occupation	Disadvantage - Household equivalised income <\$26,000 per year Advantage – Occupied private dwellings with ≥4 bedrooms
Not included*	Indigenous status, home ownership	Indigenous status	Income	Superannuation, education, occupation
Low score	Most disadvantaged – many households with low income, low education, high unemployment and many people in unskilled occupations	Most disadvantaged and least advantaged	Relatively lower education and occupation status	Relative lack of access to economic resources
High score	Most lack of disadvantage - few households with low income, low education, high unemployment and few people in unskilled occupations	Most advantaged and least disadvantaged	Relatively higher education and occupation status	Relatively greater access to economic resources

*measures of wealth (other than home ownership and income), infrastructure and health are not included in SEIFA index variables

⁵¹ Summarised from here: ABS (2023) <https://www.abs.gov.au/methodologies/socio-economic-indexes-areas-seifa-australia-methodology/2021#:~:text=The%20SEIFA%20scores%20are%20initially,averages%20of%20the%20SA1%20scores.>

Table A2-2: SEIFA data for Southern region LGAs, ranked by Index of Relative Social Disadvantage.

	IRSD		IRSAD		IER		IEO		Population
	Score	Tasmanian decile							
Brighton	855	2	862	2	924	2	849	2	18,995
Central Highlands	911	2	888	3	948	5	886	3	2,520
Clarence	1009	10	989	10	992	9	992	10	61,531
Derwent Valley	913	3	879	2	950	5	856	2	10,942
Glamorgan-Spring Bay	950	6	924	6	962	6	920	7	5,012
Glenorchy	927	4	911	5	914	2	918	6	50,411
Hobart	1046	10	1062	10	970	7	1104	10	55,077
Huon Valley	974	8	946	8	965	8	958	9	18,259
Kingborough	1038	10	1020	10	1013	10	1039	10	40,082
Sorell	991	9	946	8	993	9	932	8	16,734
Southern Midlands	951	6	911	5	1000	10	887	4	6,662
Tasman	923	4	902	4	943	4	931	7	2,593

Appendix 3: Regional economic data – income, housing, education

Table A3-1: Southern region economic data compared to Tasmania as a whole.

Key economic statistics	Southern Region	Tasmania
Median weekly household income	\$1,480	\$1,368
Individual income		
Quartile 1 income (<\$380/wk)	23.9%	25%
Quartile 2 income (\$380-701/wk)	23.3%	25%
Quartile 3 income (\$702-1,241/wk)	25.4%	25%
Quartile 4 income (>\$1,241/wk)	27.5%	25%
Equivalised household income*		
Quartile 1 income (<\$513/wk)	21.9%	25%
Quartile 2 income (\$514-881/wk)	23.9%	25%
Quartile 3 income (\$882-1,402/wk)	25.8%	25%
Quartile 4 income (>\$1,402/wk)	28.3%	25%

* calculated to be independent of household size and composition, enabling direct comparisons between datasets

Individual incomes in the Southern region are skewed towards the upper two quartiles compared to Tasmania. This is more pronounced when equivalised household income is taken into account.

Table A3-2: Housing data.

	Southern Region	Tasmania
Occupied private dwellings	88.0%	93.8%
Unoccupied private dwellings	11.7%	6.0%
Fully owed	34.6%	35.8%
Mortgage	33.1%	31.6%
Renting	26.1%	25.7%
Median weekly rent	\$348	\$290
Rental payments		
Quartile 1 (<\$212/wk)	19.9%	25%
Quartile 2 (\$212-295/wk)	16.2%	25%
Quartile 2 (\$296-387/wk)	25.7%	25%
Quartile 4 (>\$388/wk)	38.2%	25%
Housing loan repayments		
Quartile 1 (<\$920/wk)	21.6%	25%
Quartile 2 (\$920-1,357/wk)	21.7%	25%
Quartile 3 (\$1,358-1,875/wk)	26.2%	25%
Quartile 4 (>\$1,875/wk)	30.5%	25%

Housing loan data are skewed towards higher repayments per week with rental amounts very strongly so.

Table A3-3: Educational attainment data.

	Southern Region	Tasmania
Highest level of schooling – Year 9 or below	9.1%	10.8%
Highest level of schooling – Year 12	52.7%	45.5%
No qualification	36.9%	40.0%
Trade qualification	19.9%	21.9%
University qualification	27.4%	21.9%

The Southern region population is more likely than the rest of the Tasmanian population to have a higher level of educational attainment across the board, but has a much lower level of people with a trade qualification.

Table A3-4: Key employment statistics.

	Southern Region	Tasmania
Participation rate (% popn. in labour force)	60.2%	58.2%
Not in the labour force	35.0%	36.5%
Employed	93.9%	94.1%
Unemployed	6.1%	5.9%
Total labour force	145,009	270,767

The Southern region has a greater proportion of people in the labour force than Tasmania as a whole, while there is little difference in the employment/unemployment rates.



Drought Ready Tasmania

For further info contact:

Regional Drought Resilience

✉ resilience@dpac.tas.gov.au

☎ +61 3 6232 7884

Register to have your say

Scan the QR code or
visit droughtready.tas.gov.au



Australian Government
Department of Agriculture,
Fisheries and Forestry



Future
Drought
Fund



The Regional Drought Resilience Planning Program is jointly funded through the Australian Government's Future Drought Fund and the Tasmanian Government.

© 2023 State of Tasmania. Version 1 (10/2023) 230058